



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2782>

Title:	Overview of VOC emissions and chemistry from PTR-TOF-MS measurements during the SusKat-ABC campaign: high acetaldehyde, isoprene and isocyanic acid in wintertime air of the Kathmandu Valley
Authors:	Sarkar, C. (/jspui/browse?type=author&value=Sarkar%2C+C.) Sinha, V. (/jspui/browse?type=author&value=Sinha%2C+V.) Kumar, Vinod (/jspui/browse?type=author&value=Kumar%2C+Vinod)
Keywords:	VOC emissions PTR-TOF-MS SusKat-ABC campaign isocyanic acid
Issue Date:	2015
Publisher:	Copernicus GmbH
Citation:	Atmospheric Chemistry and Physics Discussions, 15(17)

Abstract: The Kathmandu Valley in Nepal suffers from severe wintertime air pollution. Volatile organic compounds (VOCs) are key constituents of air pollution, though their specific role in the Valley is poorly understood due to insufficient data. During the SusKat-ABC (Sustainable Atmosphere for the Kathmandu Valley-Atmospheric Brown Clouds) field campaign conducted in Nepal in the winter of 2012-2013, a comprehensive study was carried out to characterize the chemical composition of ambient Kathmandu air, including the determination of speciated VOCs by deploying a Proton Transfer Reaction Time of Flight Mass Spectrometer (PTR-TOF-MS)-the first such deployment in South Asia. 71 ion peaks (for which measured ambient concentrations exceeded the 2 σ detection limit) were detected in the PTR-TOF-MS mass scan data, highlighting the chemical complexity of ambient air in the Valley. Of the 71 species, 37 were found to have campaign average concentrations greater than 200 ppt and were identified based on their spectral characteristics, ambient diel profiles and correlation with specific emission tracers as a result of the high mass resolution ($m/\Delta m > 4200$) and temporal resolution (1 min) of the PTR-TOF-MS. The highest average VOC mixing ratios during the measurement period were (in rank order): acetaldehyde (8.8 ppb), methanol (7.4 ppb), acetone (4.2 ppb), benzene (2.7 ppb), toluene (1.5 ppb), isoprene (1.1 ppb), acetonitrile (1.1 ppb), C8-aromatics (~ 1 ppb), furan (~ 0.5 ppb), and C9-aromatics (0.4 ppb). Distinct diel profiles were observed for the nominal isobaric compounds isoprene (m/z Combining double low line 69.070) and furan (m/z Combining double low line 69.033). Comparison with wintertime measurements from several locations elsewhere in the world showed mixing ratios of acetaldehyde (~ 9 ppb), acetonitrile (~ 1 ppb) and isoprene (~ 1 ppb) to be among the highest reported till date. Two "new" ambient compounds namely, formamide (m/z Combining double low line 46.029) and acetamide (m/z Combining double low line 60.051), which can photochemically produce isocyanic acid in the atmosphere, are reported in this study along with nitromethane (a tracer for diesel exhaust) which has only recently been detected in ambient studies. Two distinct periods were selected during the campaign for detailed analysis: the first was associated with high wintertime emissions of biogenic isoprene, and the second with elevated levels of ambient acetonitrile, benzene and isocyanic acid from biomass burning activities. Emissions from biomass burning and biomass co-fired brick kilns were found to be the dominant sources for compounds such as propyne, propene, benzene and propanenitrile which correlated strongly with acetonitrile ($r^2 > 0.7$), a chemical tracer for biomass burning. The calculated total VOC OH reactivity was dominated by acetaldehyde (24.0 %), isoprene (20.2 %) and propene (18.7 %), while oxygenated VOCs and isoprene collectively contributed to more than 68 % of the total ozone production potential. Based on known SOA yields and measured ambient concentrations in the Kathmandu Valley, the relative SOA production potential of VOCs were: benzene > naphthalene > toluene > xylenes > monoterpenes > trimethyl-benzenes > styrene > isoprene. The first ambient measurements from any site in South Asia of compounds with significant health effects such as isocyanic acid, formamide, acetamide, naphthalene and nitromethane have been reported in this study. Our results suggest that mitigation of intense wintertime biomass burning activities, in particular point sources such as biomass co-fired brick kilns, would be important to reduce the emission and formation of toxic VOCs (such as benzene and isocyanic acid) in the Kathmandu Valley and improve its air quality.

Description: Only IISERM authors are available in the record.


URI: <https://acp.copernicus.org/preprints/15/25021/2015/acpd-15-25021-2015-supplement.pdf>
(<https://acp.copernicus.org/preprints/15/25021/2015/acpd-15-25021-2015-supplement.pdf>)
<http://hdl.handle.net/123456789/2782> (<http://hdl.handle.net/123456789/2782>)

Appears in Research Articles (</jspui/handle/123456789/9>)
Collections:

Files in This Item:

File	Description	Size	Format	
Need to add pdf.odt (/jspui/bitstream/123456789/2782/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text	View/Open (/jspui/bitstream/123456789/2782/1/Need%20to%20add%20pdf.odt)

Show full item record (</jspui/handle/123456789/2782?mode=full>)

 (</jspui/handle/123456789/2782/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.