



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)


Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2727>

Title:	Direct Bis-Arylation of Cyclobutanecarboxamide via Double C–H Activation: An Auxiliary-Aided Diastereoselective Pd-Catalyzed Access to Trisubstituted Cyclobutane Scaffolds Having Three Contiguous Stereocenters and an All-cis Stereochemistry
Authors:	Parella, R. (/jspui/browse?type=author&value=Parella%2C+R.) Gopalakrishnan, B. (/jspui/browse?type=author&value=Gopalakrishnan%2C+B.) Babu, S.A. (/jspui/browse?type=author&value=Babu%2C+S.A.)
Keywords:	Double C–H activation Methylene Pd-catalyzed Auxiliary
Issue Date:	2013
Publisher:	American Chemical Society
Citation:	Journal of Organic Chemistry, 78(23), pp.11911-11934.
Abstract:	An auxiliary-aided Pd-catalyzed highly diastereoselective double C–H activation and direct bis-arylation of methylene C(sp ³)–H bonds of cyclobutanecarboxamides and the syntheses of several novel trisubstituted cyclobutanecarboxamide scaffolds having an all-cis stereochemistry are reported. Extensive screening of various auxiliaries and reaction conditions was performed to firmly establish the optimized reaction conditions required for effecting the mono- or double C–H arylation of cyclobutanecarboxamides. The auxiliary-attached cyclobutanecarboxamides 15a, 15g, and 15h, prepared from the auxiliaries such as, 8-aminoquinoline, 2-(methylthio)aniline, and N',N'-dimethylethane-1,2-diamine were found to undergo an efficient direct bis-arylation. The Pd-catalyzed arylation reaction of N-(quinolin-8-yl)cyclobutanecarboxamide 15a with one equivalent or more of aryl iodides, afforded the corresponding bis-arylated cyclobutanecarboxamides 16a–y. Nevertheless, the Pd-catalyzed arylation of 15a with just 0.5 equiv of the aryl iodides 13a, 13b, 13e, and 13m, selectively gave the corresponding monoarylated cyclobutanecarboxamides 17a–17d. The Pd-catalyzed arylation of 15g or 15h with one equivalent or more of aryl iodides afforded the bis-arylated cyclobutanecarboxamides 19a–19c and 21a–21m, respectively. However, the Pd-catalyzed arylations of compounds 15g or 15h with just 0.5 equiv of aryl iodides were ineffective. The stereochemistry of compounds obtained in this work was unambiguously assigned from the X-ray structures of representative products.
URI:	https://pubs.acs.org/doi/10.1021/jo4019733 (https://pubs.acs.org/doi/10.1021/jo4019733) http://hdl.handle.net/123456789/2727 (http://hdl.handle.net/123456789/2727)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format	
Need to add pdf.odt (/jspui/bitstream/123456789/2727/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text	View/Open (/jspui/bitstream/123456789/2727/1/Need%20to%20add%20pdf.odt)

Show full item record (</jspui/handle/123456789/2727?mode=full>)

 (</jspui/handle/123456789/2727/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.