



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Thesis & Dissertation (/jspui/handle/123456789/1)
/ Master of Science (/jspui/handle/123456789/2)
/ MS-11 (/jspui/handle/123456789/537)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/589>

Title: Aqueous Phase Oxidation of SO₂ During Winter Fog at a Sub Urban Site in North-West Indo Gangetic Plain

Authors: Vaishali (/jspui/browse?type=author&value=Vaishali)

Keywords: Chemistry
Atmospheric Chemistry
Environmental Science
Sulfur dioxide
Air Pollution

Issue Date: 9-Aug-2016

Publisher: IISER-M

Abstract: Sulfur dioxide is an important air pollutant that is emitted into the air from anthropogenic and natural sources. High SO₂ mixing ratios are known for their serious health implications, as the gas is oxidized to sulfuric acid within the human lung. Even at much lower mixing ratios SO₂ can affect forest ecosystems negatively. The gas is efficiently accumulated on leaves (dry deposition) and acid is formed in situ in the presence of water, which results in acid rain. Due to these environmental consequences, SO₂ emissions are increasingly regulated and have decreased since the 1990s in most countries around the world. It is important to study and understand the global sulfur cycle primarily to the large impact of sulfate aerosol formed on the global aerosol radiative forcing -0.41 ± 0.2 W/m², which currently offsets between 10-36% of the CO₂ induced warming. The 50% uncertainty in this global forcing estimate of anthropogenic SO₂ emissions, is largely due to uncertainties associated with the treatment of the SO₂ oxidation to sulfate in global climate models. In the study I will discuss important empirical observations pertaining to the aqueous phase oxidation of SO₂ in wintertime fog in the NW-IGP. The study looks at visibility reduction due to presence of fog and classifies days with a visibility below 2 km throughout as foggy. For those foggy days, I investigate the changes of SO₂ mixing ratios that coincide with simultaneous changes in the aerosol liquid water content to quantify the effective reactive uptake of SO₂ (g) and NH₃ (g) into the aerosol liquid phase. I have found that at our site SO₂ is fully neutralized and has been converted to (NH₄)₂SO₄. Therefore I have predicted that the fog water pH should be close to neutral which well correlates with the experimental pH measured during winter 2015-16. Out of 12 global models, 4 models fix pH in the Northern Hemisphere and the others treat it with an equation valid from pH 3.5-5.5. Five models participate in IPCC-AR5 but no model is capable of treating aqueous phase chemistry of SO₂ when pH of the system goes beyond 5. As India and China contribute to 40% of the total global SO₂ budget, it becomes more important to include all sinks of SO₂ to estimate correct budget. These are densely populated areas with large portion of land use being agricultural thus having high ammonia emissions from this sector which affects the lifetime and forcing of sulfate. Thus amount of aqueous phase sulfate oxidation is underestimated which needs to be taken care of, especially when the systems are neutral as it would help to improve the global budget of SO₂ and total radiative forcing due to sulfate.


URI: <http://hdl.handle.net/123456789/589> (<http://hdl.handle.net/123456789/589>)

Appears in Collections: MS-11 (/jspui/handle/123456789/537)

Files in This Item:

File	Description	Size	Format	
MS Thesis Vaishali MS11074.pdf (/jspui/bitstream/123456789/589/3/MS%20Thesis%20Vaishali%20MS11074.pdf)		1.91 MB	Adobe PDF	View/Open (/jspui/bitstream/123456789/589/3/MS%20Thesis%20Vaishali%20MS11074.pdf)

[Show full item record \(/jspui/handle/123456789/589?mode=full\)](/jspui/handle/123456789/589?mode=full)

 [\(/jspui/handle/123456789/589/statistics\)](/jspui/handle/123456789/589/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.