



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4934>


| | |
|-------------------------|--|
| Title: | Size dependence of CdS nanoparticles on the precursor concentration and visible light driven photocatalytic degradation of methylene blue |
| Authors: | Thakur, Smriti (/jspui/browse?type=author&value=Thakur%2C+Smriti) Kaur, Rupinder (/jspui/browse?type=author&value=Kaur%2C+Rupinder) Mandal, Sanjay K. (/jspui/browse?type=author&value=Mandal%2C+Sanjay+K.) |
| Keywords: | Size dependence of CdS nanoparticles photocatalytic methylene blue |
| Issue Date: | 2021 |
| Publisher: | New Journal of Chemistry |
| Citation: | New Journal of Chemistry, 45(27), 12227–12235. |
| Abstract: | Using a Cd(II) coordination polymer, [Cd2(bpma)2(adc)2] _n (where bpma = N,N'-bis(pyridylmethyl) amine and adc = acetylene dicarboxylate), as the source of Cd ²⁺ ions and thiourea as the source of S ²⁻ ions in various ratios (1 : 3, 1 : 5, 1 : 7, 1 : 10, 1 : 15 and 1 : 20) in water, CdS nanoparticles (CdS-1 to CdS-6) were synthesized under hydrothermal conditions without a template and an added surfactant. The size of nanoparticles increases with an increase in the concentration of thiourea. These nanoparticles were characterized by powder X-ray diffraction (XRD), field emission scanning electron microscopy (FESEM), energy dispersive X-ray spectroscopy (EDS), transmission electron microscopy (TEM), and UV–vis diffuse reflectance spectroscopy. The XRD results revealed the presence of both cubic and hexagonal forms in CdS-1 to CdS-6. Based on their UV–vis diffuse reflectance, the band gap energies were estimated to be 2.26, 2.20, 2.19, 2.18, 2.17 and 2.15 eV for CdS-1, CdS-2, CdS-3, CdS-4, CdS-5 and CdS-6, respectively. The photoluminescence spectra of CdS-1 to CdS-6 (excitation wavelength: 350 nm) displayed a blue emission in the wavelength range of 434–437 nm and a green emission in the range of 533–552 nm. The blue emission resulted from the recombination of excitons, whereas the green emission was caused by sulphur vacancies in CdS nanoparticles. Furthermore, the photocatalytic degradation of the methylene blue dye was carried out in water under visible light using CdS-1 to CdS-4, where the maximum degradation efficiency of 97% was obtained using 0.02 mol% of CdS-1. |
| Description: | Only IISERM authors are available in the record. |
| URI: | https://doi.org/10.1039/d1nj01588e (https://doi.org/10.1039/d1nj01588e) http://hdl.handle.net/123456789/4934 (http://hdl.handle.net/123456789/4934) |
| Appears in Collections: | Research Articles (/jspui/handle/123456789/9) |

Files in This Item:

| File | Description | Size | Format |
|------|-------------|------|--------|
| | | | |

| | | | | |
|--|--|-------------|---------|------------------------------------|
| Need To Add...Full Text_PDF. (/jspui/bitstream/123456789/4934/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF.) | Only IISERM authors are available in the record. | 15.36 kB | Unknown | View/Open (/jspui/ |
|--|--|-------------|---------|------------------------------------|

[Show full item record \(/jspui/handle/123456789/4934?mode=full\)](#)

 [\(/jspui/handle/123456789/4934/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.