



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**

**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1739>


Title:	How different is the borazine–acetylene dimer from the benzene–acetylene dimer? A matrix isolation infrared and ab initio quantum chemical study
Authors:	Verma, Kanupriya (/jspui/browse?type=author&value=Verma%2C+Kanupriya) Sathyamurthy, N. (/jspui/browse?type=author&value=Sathyamurthy%2C+N.) Viswanathan, K.S. (/jspui/browse?type=author&value=Viswanathan%2C+K.S.)
Keywords:	Borazine acetylene hydrogen bonded interaction $\pi$ -stacked interaction matrix isolation infrared spectroscopy
Issue Date:	2017
Publisher:	T&F
Citation:	Molecular Physics, 115 (21-22)
Abstract:	The 1:1 dimer of borazine–acetylene has been studied for the first time, both experimentally and computationally. The borazine–acetylene dimer was trapped in Ar and N <sub>2</sub> matrices, and studied using infrared spectroscopy. Our experiments clearly revealed two isomers of the borazine–acetylene complex, one in which the N–H of borazine interacted with the carbon of acetylene, and another in which the C–H of acetylene formed a hydrogen bond with a nitrogen atom of borazine. The formation of both isomers in the matrix was evidenced by shifts in the vibrational frequencies of the appropriate modes. Reassuringly, the experimental observations were corroborated by our computations using the second-order Møller–Plesset perturbation theoretic method and coupled-cluster singles, doubles and perturbative triples method in conjunction with different Dunning basis sets, which indicated both these isomers to be stable minima, with the N–H...C complex being the global minimum. Atoms-in-molecules and energy decomposition analysis were also carried out for the different isomers of the dimer. These studies reveal that replacing the three C–C linkages in benzene with three B–N linkages in borazine modifies the interaction in the dimer sufficiently, to result in a different potential energy landscape for the borazine–acetylene system when compared with the benzene–acetylene system.
Description:	Only IISERM authors are available in the record
URI:	<a href="https://www.tandfonline.com/doi/full/10.1080/00268976.2017.1284357">https://www.tandfonline.com/doi/full/10.1080/00268976.2017.1284357</a> ( <a href="https://www.tandfonline.com/doi/full/10.1080/00268976.2017.1284357">https://www.tandfonline.com/doi/full/10.1080/00268976.2017.1284357</a> ) <a href="http://hdl.handle.net/123456789/1739">http://hdl.handle.net/123456789/1739</a> ( <a href="http://hdl.handle.net/123456789/1739">http://hdl.handle.net/123456789/1739</a> )
Appears in	Research Articles (/jspui/handle/123456789/9)
Collections:	

Files in This Item:

File	Description	Size	Format
Need to add pdf.ott (/jspui/bitstream/123456789/1739/1/Need%20to%20add%20pdf.ott)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/1739/1/Need%20to%20add%20pdf.ott\)](#)

Show full item record (</jspui/handle/123456789/1739?mode=full>)

 (</jspui/handle/123456789/1739/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.