

Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

- / Publications of IISER Mohali (/jspui/handle/123456789/4)
- / Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/3966

Title: Spectral States of OJ 287 blazar from Multiwavelength Observations with AstroSat

Authors: Singh, K.P. (/jspui/browse?type=author&value=Singh%2C+K.P.)

Keywords: Radiation mechanisms: non-thermal

Galaxies: active Galaxies: jets Gamma-rays: galaxies X-rays: galaxies

Issue Date: 2021

Publisher: Oxford Academic

Citation: Monthly Notices of the Royal Astronomical Society, Volume 509(2):2696–2706.

Abstract:

We present AstroSat soft X-ray, near-UV (NUV), and far-UV (FUV) observations of a blazar, OJ 287, carried out in 2017, 2018, and 2020. The simultaneous observations with NuSTAR in 2017 provide a broad-band look encompassing NUV, FUV, soft, and hard X-rays. Captured in three different broad-band spectral states in three observations, the X-ray spectrum is found to be the hardest during 2018, while the high-energy-end of the simultaneous optical-FUV spectrum shows a steepening that is modelled with a broken power-law spectrum. The spectral energy distribution (SED) in 2017 shows a relatively flatter optical-FUV and soft X-ray spectra, implying an additional emission component. The 2020 optical-FUV spectrum is harder than in 2017 and 2018, with an extremely soft X-ray spectrum and a hardening above \sim 1 GeV, similar to the SEDs of Highenergy-peaked BL Lac objects (HBL), thereby establishing that this additional emission component has HBL-like properties. The AstroSat multiwavelength observations trace the spectral evolution from the end-phase of the HBL component in 2017 to its disappearance in 2018 followed by its revival in 2020. A single zone leptonic model reproduces the 2018 broad-band spectrum, while the 2017 and 2020 SEDs require an additional HBL-like emitting zone. The spectral evolution of the high-energy-end of optical-UV spectrum, revealed by the FUV observations in 2017 and 2018, strongly suggests that X-ray spectral changes in the normal broad-band spectral state of OJ 287 are primarily due to the evolution of the optical-UV synchrotron spectrum.

Description: Only IISERM authors are available in the record.

URI: https://academic.oup.com/mnras/article/509/2/2696/6415211

(https://doi.org/10.1093/mpros/stab3161/https://doi.org/1

https://doi.org/10.1093/mnras/stab3161 (https://doi.org/10.1093/mnras/stab3161) http://hdl.handle.net/123456789/3966 (http://hdl.handle.net/123456789/3966)

Appears in Collections:

Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File Description Size Format

Need to add pdf.docx
(/jspui/bitstream/123456789/3966/1/Need%20to%20add%20pdf.docx)

Need to add pdf.docx
(/jspui/bitstream/123456789/3966/1/Need%20to%20add%20pdf.docx)

View/Open (/jspui/bitstream/123456789/3966/1/Need%20to%20add%20pdf.docx)

Show full item record (/jspui/handle/123456789/3966?mode=full)

. I (/jspui/handle/123456789/3966/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.