



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Thesis & Dissertation (/jspui/handle/123456789/1)

/ Doctor of Philosophy (PhD) (/jspui/handle/123456789/268)

/ PhD-2014 (/jspui/handle/123456789/462)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1557>

Title:	Scanning Tunnelling Microscopy and Transport Spectroscopy on Candidate Topological Systems
Authors:	Das, Shekhar (/jspui/browse?type=author&value=Das%2C+Shekhar)
Keywords:	Scanning technique in STM Topological invariant Topological superconductors Superconductivity in PdTe ₂
Issue Date:	Oct-2019
Publisher:	IISER Mohali

Abstract: Topologically non-trivial systems have emerged as a new area of research in condensed matter physics. The study of topologically nontrivial electronic systems is important because they provide a platform to experimentally investigate new low-energy excitations. In this thesis, I will discuss three such systems: Conventional Superconductivity in PdTe₂: PdTe₂ is a unique topological system where type II Dirac semimetallic phase coexists with a superconducting phase. This combined nature offers an interesting material candidate for investigation of possible topological superconductivity. We have performed high-resolution scanning tunnelling spectroscopic (STS) measurements to study the nature of superconductivity in this system. Theoretically, it has been speculated that superconductivity in such systems will be unconventional in nature. However, our experiments provide direct evidence of conventional superconductivity in PdTe₂. High spin-polarization in Sb₂Se₃: Most of the A₂B₃ type chalcogenides (e.g. Bi₂Se₃, Bi₂Te₃ and Sb₂Te₃ etc.) are well known topological insulators. Though Sb₂Se₃ is a member of the same group, it is a band insulator under ambient conditions. A prominent quasiparticle interference (QPI) pattern observed in STM conductance mapping indicates the presence of backscattering, which is forbidden for a topological insulator. Interestingly, like most of the topological insulators, Sb₂Se₃ shows a high spin polarization (69 %) in transport current. To understand the origin of such high spin polarization, we performed band structure calculations which revealed presence of two trivial surface states with one undergoing large splitting due to Rashba type SOC and leading to high spin polarization. Tip-induced superconductivity in Pb_{0.6}Sn_{0.4}Te: The emergence of tip-induced superconductivity (TISC) has offered a possible way to detect topological superconductivity induced by a point contacts. We performed similar experiment on a topological crystalline insulator (TCI) Pb_{0.6}Sn_{0.4}Te with non-superconducting tips. Our experiments revealed that Ag/Pb_{0.6}Sn_{0.4}Te and Pd/Pb_{0.6}Sn_{0.4}Te point-contacts behave as a superconductor. As expected for a superconductor, we observed a sharp resistive transition at 6 K which evolved systematically with an applied magnetic field. This is also confirmed through temperature and magnetic field dependent studies of the point contact spectra. Moreover, on the basis of H_c Vs T_c phase diagram we predict that the induced superconductivity is conventional in nature. Furthermore, I will discuss low temperature and high vacuum point contact probe designing and fabrication.

URI: <http://hdl.handle.net/123456789/1557> (<http://hdl.handle.net/123456789/1557>)


Appears in PhD-2014 (</jspui/handle/123456789/462>)

Collections:

Files in This Item:

File	Description	Size	Format	
PH14012.pdf (/jspui/bitstream/123456789/1557/3/PH14012.pdf)		8.9 MB	Adobe PDF	View/Open (/jspui/bitstream/123456789/1557/3/)

[Show full item record \(/jspui/handle/123456789/1557?mode=full\)](/jspui/handle/123456789/1557?mode=full)

 (</jspui/handle/123456789/1557/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

Admin Tools

[Edit...](#)

[Export Item](#)

[Export \(migrate\) Item](#)

[Export metadata](#)

