



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1809>


Title:	Synthesis, characterizations and single crystal structure of di-nuclear azido-bridged Cd(II) coordination polymer with Schiff base precursor (H ₂ LpentOMe): DFT, fluorescence, solvatochromism and in vitro antimicrobial assay
Authors:	Das, Dhiraj (/jspui/browse?type=author&value=Das%2C+Dhiraj)
Keywords:	Azide Antibacterial Cd(II) DFT Schiff base
Issue Date:	2019
Publisher:	Elsevier
Citation:	Inorganica Chimica Acta, 496.
Abstract:	<p>A di-nuclear azido-bridged Cd(II) 1-D coordination polymer [Cd₂(H₂LpentOMe)(μ_{1,1}-N₃)₂]_n (1) has been successfully synthesized using less explored bi-compartmental Schiff base ligand (H₂LpentOMe) and characterized by elemental analysis, FT-IR, FT-Raman, UV-Visible, SEM-EDAX, powder X-ray diffraction and fluorescence spectroscopy. Solid-state X-ray single crystal study revealed two different geometrical environment of Cd metal centres with distorted square pyramidal (Cd1) and distorted pentagonal bipyramid (Cd2). Overall, small-sized azide ions in asymmetric unit act as μ_{1,1} bridging mode. Geometry is optimized in gas phase using ORCA 3.0.3 and B3LYP level TZVP basis set to explain frontier molecular orbitals (FMO), molecular electrostatic potential (MEP) and global reactivity. The HOMO-LUMO energy gap (3.434 eV) suggests that chemical reactivity of complex 1 is low but fairly stable. Moreover, molecular electrostatic potential map was drawn to identify reactive regions in terms of electrophilic and nucleophilic. The steady state and time-resolved fluorescence properties have been explored in DCM and solid-state condition at room temperature. Complex 1 exhibit bi and tri-exponential decay in DCM as well as solid-state. The fluorescence behaviours are predominantly intra-ligand in nature (π → π*) with lifetimes in the range (1.16–1.11 ns). Solvatochromism has been reported to show solvent dependent absorption and fluorescence spectral changes. Fascinatingly, red shifted solvatochromism was observed upon increasing solvent polarity. Finally, concentration dependent antimicrobial activity was investigated against two standard bacterial strains (Staphylococcus aureus (ATCC 25923) and Methicillin-resistant Staphylococcus aureus (MRSA) where 250 μg/ml concentration of complex 1 is sufficient to show the antimicrobial effect.</p>
Description:	Only IISERM authors are available in the record.
URI:	https://www.sciencedirect.com/science/article/pii/S0020169319308485 (https://www.sciencedirect.com/science/article/pii/S0020169319308485) http://hdl.handle.net/123456789/1809 (http://hdl.handle.net/123456789/1809)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
------	-------------	------	--------

Need to add pdf.odt (/jspui/bitstream/123456789/1809/1/Need%20to%20add%20pdf.odt)	8.63 kB	OpenDocument Text	View/Open (/jspui/bitstream/123456789/1809/1/Need%20to%20add%20pdf.odt)
--	------------	----------------------	---

Show full item record (/jspui/handle/123456789/1809?mode=full)

 (/jspui/handle/123456789/1809/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.