



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/3254>

Title: Vegetational responses to monsoon variability during Late Holocene: Inferences based on carbon isotope and pollen record from the sedimentary sequence in Dzukou valley, NE India

Authors: Bhattacharya, S. (/jspui/browse?type=author&value=Bhattacharya%2C+S.)
Ambili, Anoop (/jspui/browse?type=author&value=Ambili%2C+Anoop)

Keywords: Biomarkers
Carbon isotope
Late Holocene
Monsoon
NE India
Pollens

Issue Date: 2020

Publisher: Elsevier B.V

Citation: Catena, 194

Abstract: The study aims to identify regional vegetational changes during the late Holocene from the Dzukou valley in north-eastern (NE) India and the potential relationship of these vegetation dynamics with climatic shifts. The investigations utilize a combination of $\delta^{13}\text{C}_{\text{org}}$ of organic matter, lipid biomarker (n-alkane) as well as pollen data derived from ~1.4 m sedimentary profile spanning the last ~3150 cal BP. The $\delta^{13}\text{C}_{\text{org}}$ values during late Holocene range from -29.1‰ to -27.7‰ suggesting that organic carbon was derived exclusively from C3 vegetation. The profile is characterized by a homologous series of n-alkanes ranging in chain length from n-C18 to n-C35 with maxima at n-C29. The generated proxy data reveal three phases of climatic and vegetation shifts with the first phase from ~3150 cal BP to ~2300 cal BP being characterised by relatively high $\delta^{13}\text{C}_{\text{org}}$ values and occupied by Pine-Oak forest indicating a comparatively dry climatic conditions. Subsequently, the second phase from ~2300 cal BP to ~1000 cal BP shows an increase in arboreal pollen (tree elements) and gradually decreasing trend of $\delta^{13}\text{C}_{\text{org}}$ with a total shift of ~1.4‰ suggesting moist climatic conditions. During the third phase from ~1000 cal BP onwards, the precipitation declined till date as suggested by increasing trend in $\delta^{13}\text{C}_{\text{org}}$ values and good recovery of Pinus-Oak forest pollens. This study provides significant insights into the use of $\delta^{13}\text{C}_{\text{org}}$ and biomarkers along with pollen data to understand the influence of fluctuating rainfall (in a high rainfall zone) in altering the floristic composition within a region. Thus, the study has provided ground for further investigations to address issues related to biodiversity loss, extent of endemism in high rainfall zones dominated by single types (C3 and/or C4) of vegetation

Description: Only IISERM authors are available in the record.

URI: <https://www.sciencedirect.com/science/article/abs/pii/S0341816220302472>
(<https://www.sciencedirect.com/science/article/abs/pii/S0341816220302472>)
<http://hdl.handle.net/123456789/3254> (<http://hdl.handle.net/123456789/3254>)

Appears in Collections: Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
------	-------------	------	--------

Need to add pdf.odt (/jspui/bitstream/123456789/3254/1/Need%20to%20add%20pdf.odt)	8.63 kB	OpenDocument Text	View/Open (/jspui/bitstream/123456789/3254/1/Need%20to%20add%20pdf.odt)
--	------------	----------------------	---

Show full item record (/jspui/handle/123456789/3254?mode=full)

 (/jspui/handle/123456789/3254/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.