



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4343>


Title:	On the understanding of surface ozone variability, its precursors and their associations with atmospheric conditions over the Delhi region
Authors:	Attada, Raju (/jspui/browse?type=author&value=Attada%2C+Raju)
Keywords:	Annual cycle High ozone days Photochemical reactions Surface ozone
Issue Date:	2021
Publisher:	Elsevier
Citation:	Atmospheric Research, 258, 105653.
Abstract:	<p>The present study investigates the variability of surface ozone (O₃) and its precursors (NO_x, CO) at different time scales for 3-year period (2016–2018) using continuous ground-based observations from 25 stations located within the Indian Capital Delhi (28.83°N, 77.81°E). Observations indicate that there is a clear spatial heterogeneity and seasonality in surface O₃ where larger magnitudes (40–60 ppb) are observed during pre-monsoon season, and 10–20 ppb in other seasons. The frequency distribution exhibits that 70% of O₃ concentrations are in the range of 10–50 ppb, while 10% lies larger than 50 ppb and the remaining 20% is mostly confined within 50–80 ppb. Further, a detailed analysis on the diurnal cycle of surface ozone and its precursors performed over IMD Lodhi Road location. Analysis of diurnal cycles show maximum and minimum O₃ concentrations during the morning and nocturnal hours, with a rate of change 4.8 ± 1.96 and -7.4 ± 0.95 ppb h⁻¹, respectively. On the other hand, the rate of change of O₃ during high ozone days (HOD) is found to be 5.61 ± 2.6 ppb h⁻¹ in the morning and -8.04 ± 3.4 ppb h⁻¹ in the evening hours, respectively. It is also noted that the increase in precursor concentrations and associated titration effects appear to play a considerable role in surface O₃ depletion during the evening hours. It is also noted from the composites of HODs that there is an enhanced diurnal O₃ (80–90 ppb) concentrations compared to 3-year average (40–60 ppb). Further evaluation of associations between surface O₃ changes and meteorological parameters shows that observed surface O₃ (Copernicus Atmosphere Monitoring Service-CAMS modelled O₃) show positive correlations of $R^2 = 0.3$ (0.42) with temperature, of $R^2 = 0.47$ (0.60) with Solar Radiation (SR), of $R^2 = 0.39$ (0.59) with boundary layer height. An overall assessment shows CAMS modelled three year daily averaged ozone values are generally in good agreement with surface ozone observations with a CC value of about $R^2 = 0.65$.</p>
Description:	Only IISER Mohali authors are available in the record.
URI:	https://doi.org/10.1016/j.atmosres.2021.105653 (https://doi.org/10.1016/j.atmosres.2021.105653) http://hdl.handle.net/123456789/4343 (http://hdl.handle.net/123456789/4343)
Appears in	Research Articles (/jspui/handle/123456789/9)
Collections:	

Files in This Item:

File	Description	Size	Format
------	-------------	------	--------

Need To Add...Full Text_PDF..pdf (/jspui/bitstream/123456789/4343/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF..pdf)	Only IISER Mohali authors are available in the record.	15.36 kB	Adobe PDF	View/Open (/jspu
--	--	-------------	--------------	----------------------------------

[Show full item record \(/jspui/handle/123456789/4343?mode=full\)](/jspui/handle/123456789/4343?mode=full)

 [\(/jspui/handle/123456789/4343/statistics\)](/jspui/handle/123456789/4343/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.