



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1718>


| | |
|--------------|--|
| Title: | Dissipative properties and isothermal compressibility of hot and dense hadron gas using non-extensive statistics |
| Authors: | Tiwari, Satish Kumar (/jspui/browse?type=author&value=Tiwari%2C+Satish+Kumar) Kakati, N. (/jspui/browse?type=author&value=Kakati%2C+N.) |
| Keywords: | hadron gas isothermal compressibility non-extensive statistics hot and dense |
| Issue Date: | 2018 |
| Publisher: | Springer New York LLC |
| Citation: | European Physical Journal C, 78(11). |
| Abstract: | We evaluate the transport properties such as shear viscosity (η), bulk viscosity (ζ) and their ratios over entropy density (s) for hadronic matter using relativistic non-extensive Boltzmann transport equation (NBTE) in relaxation time approximation (RTA). In NBTE, we argue that the system far from equilibrium may not reach to an equilibrium described by extensive (Boltzmann-Gibbs (BG)) statistics but to a q-equilibrium defined by Tsallis non-extensive statistics after subsequent evolution, where q denotes the degree of non-extensivity. We observe that η/s and ζ/s decrease rapidly with temperature (T) for various q-values. As q increases, the magnitudes of η/s and ζ/s decrease with T. We also show the upper mass cutoff dependence of these ratios for a particular q and find that they decrease with the increase in mass cutoff of hadrons. Further, we present the first estimation of isothermal compressibility (κT) using non-extensive Tsallis statistics at finite baryon chemical potential (μB). It is observed that, κT changes significantly with the degree of non-extensivity. We also study the squared speed of sound (c^2s) as a function of temperature at finite baryon chemical potential for various q and upper mass cutoffs. It is noticed that there is a strong impact of q and mass cutoff on the behaviour of c^2s . |
| Description: | Only IISERM authors are available in the record. |
| URI: | https://link.springer.com/article/10.1140/epjc/s10052-018-6411-y (https://link.springer.com/article/10.1140/epjc/s10052-018-6411-y) http://hdl.handle.net/123456789/1718 (http://hdl.handle.net/123456789/1718) |
| Appears in | Research Articles (/jspui/handle/123456789/9) |
| Collections: | |

Files in This Item:

| File | Description | Size | Format |
|--|-------------|------------|----------------------|
| Need to add pdf.odt (/jspui/bitstream/123456789/1718/1/Need%20to%20add%20pdf.odt) | | 8.04 kB | OpenDocument Text |

[View/Open \(/jspui/bitstream/123456789/1718/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/1718?mode=full>)

 (</jspui/handle/123456789/1718/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.