



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4438>


Title:	N-Terminal Extensions Appear to Frustrate HU Heterodimer Formation by Strengthening Intersubunit Contacts and Blocking the Formation of a Heterotetrameric Intermediate
Authors:	Arora, Kanika (/jspui/browse?type=author&value=Arora%2C+Kanika) Thakur, Bhishem (/jspui/browse?type=author&value=Thakur%2C+Bhishem) Mrigwani, Arpita (/jspui/browse?type=author&value=Mrigwani%2C+Arpita) Guptasarma, Purnananda (/jspui/browse?type=author&value=Guptasarma%2C+Purnananda)
Keywords:	HU Heterodimer Strengthening Intersubunit Heterotetrameric N-Terminal
Issue Date:	2021
Publisher:	ACS Publication
Citation:	Biochemistry, 60(23), 1836–1852.
Abstract:	<p>HU is a bacterial nucleoid-associated protein. Two homologues, known as HU-A, and HU-B, are found in <i>Escherichia coli</i> within which the early, late, and stationary phases of growth are dominated by HU-AA, HU-BB, and HU-AB dimers, respectively. Here, using genetic manipulation, mass spectrometry, spectroscopy, chromatography, and electrophoretic examination of glutaraldehyde-mediated cross-linking of subunits, in combination with experiments involving mixing, co-expression, unfolding, and refolding of HU chains, we show that the spontaneous formation of HU-AB heterodimers that is reported to occur upon mixing of wild-type HU-AA and HU-BB homodimers does not occur if chains possess N-terminal extensions. We show that N-terminal extensions interfere with the conversion of homodimers into heterodimers. We also show that heterodimers are readily formed at anticipated levels by chains possessing N-terminal extensions in vivo, when direct chain–chain interactions are facilitated through production of HU-A and HU-B chains from proximal genes located upon the same plasmid. From the data, two explanations emerge regarding the mechanism by which N-terminal extensions happen to adversely affect the conversion of homodimers into heterodimers. (1) The disappearance of the α-amino group at HU's N-terminus impacts the intersubunit stacking of β-sheets at HU's dimeric interface, reducing the ease with which subunits dissociate from each other. Simultaneously, (2) the presence of an N-terminal extension appears to sterically prevent the association of HU-AA and HU-BB homodimers into a critically required, heterotetrameric intermediate (within which homodimers could otherwise exchange subunits without releasing monomers into solution, by remaining physically associated with each other).</p>
Description:	Only IISER Mohali authors are available in the record.
URI:	https://doi.org/10.1021/acs.biochem.1c00081 (https://doi.org/10.1021/acs.biochem.1c00081) http://hdl.handle.net/123456789/4438 (http://hdl.handle.net/123456789/4438)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
------	-------------	------	--------

Need To Add...Full Text_PDF..pdf (/jspui/bitstream/123456789/4438/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF..pdf)	Only IISER Mohali authors are available in the record.	15.36 kB	Adobe PDF	View/Open (/jspu
--	--	-------------	--------------	----------------------------------

[Show full item record \(/jspui/handle/123456789/4438?mode=full\)](#)

 [\(/jspui/handle/123456789/4438/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.