



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Thesis & Dissertation (/jspui/handle/123456789/1)

/ Doctor of Philosophy (PhD) (/jspui/handle/123456789/268)

/ PhD-2008 (/jspui/handle/123456789/1096)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/625>


Title:	Characterizing Quantum Correlations in the Nonsignaling Framework
Authors:	Jebarathinam, C. (/jspui/browse?type=author&value=Jebarathinam%2C+C.)
Keywords:	Physics Quantum Theory
Issue Date:	4-Sep-2016
Publisher:	IISER-M
Abstract:	<p>Bell nonlocality of quantum theory refers to the nonclassical correlations obtained by local measurements on spatially separated entangled subsystems. Bell nonlocality is a resource for device-independent quantum information processing. Quantum discord was introduced as a measure of quantum correlations which captures nonclassical correlations in separable states as well. Recently, it has been shown that non-null quantum discord is a resource for quantum information processing. Quantum correlations forms a subset of the set of nonsignaling boxes. This allows us to characterize quantum correlations as a convex combination of the extremal boxes of the nonsignaling polytope which are Popescu-Rohrlich boxes (maximally nonlocal boxes) and local deterministic boxes. There exists multiple decomposition of quantum correlations in the context of the nonsignaling polytope. I find that the existence of Popescu-Rohrlich box decomposition for local boxes associates two notions of discord which capture nonclassicality of quantum correlations originating from Bell nonlocality and EPR-steering. I introduce, Bell and Mermin discord, and show that any bipartite nonsignaling box admits a three-way decomposition. This decomposition allows us to isolate the origin of nonclassicality into three disjoint sources: a Popescu-Rohrlich box, a maximally EPR-steerable box, and a classical correlation. Interestingly, I show that all non-null quantum discord states which are neither classical-quantum states nor quantum-classical states can give rise to nonclassical correlations which have non-null Bell and/or Mermin discord for suitable incompatible measurements. I introduce two notions of genuine discord, which are the generalizations of Bell and Mermin discord to the multipartite scenario, to characterize the presence of genuine nonclassicality in quantum correlations.</p>
URI:	http://hdl.handle.net/123456789/625 (http://hdl.handle.net/123456789/625)

Appears in PhD-2008 (/jspui/handle/123456789/1096)
Collections:

Files in This Item:

File	Description	Size	Format	
PH-08003.pdf (/jspui/bitstream/123456789/625/1/PH-08003.pdf)		1.29 MB	Adobe PDF	View/Open (/jspui/bitstream/123456789/625/1/PH-08003.pdf)

[Show full item record \(/jspui/handle/123456789/625?mode=full\)](#)

 [\(/jspui/handle/123456789/625/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

Admin Tools

[Edit...](#)

[Export Item](#)

[Export \(migrate\) Item](#)

[Export metadata](#)