



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**

**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1627>


Title:	Intermolecular Charge-Transfer Modulates Liquid-Liquid Phase Separation and Liquid-to-Solid Maturation of an Intrinsically Disordered pH-Responsive Domain
Authors:	Dogra, P. (/jspui/browse?type=author&value=Dogra%2C+P.) Joshi, Ashish (/jspui/browse?type=author&value=Joshi%2C+Ashish) Majumdar, A. (/jspui/browse?type=author&value=Majumdar%2C+A.) Mukhopadhyay, S. (/jspui/browse?type=author&value=Mukhopadhyay%2C+S.)
Keywords:	Charge transfer Peptides and proteins Liquids Fluorescence
Issue Date:	2019
Publisher:	ACS Publications
Citation:	Journal of the American Chemical Society, 141(51), pp. 20380-20389.
Abstract:	Liquid-liquid phase separation of intrinsically disordered proteins into mesoscopic, dynamic, liquid-like supramolecular condensates is thought to govern critical cellular functions. These condensates can mature from a functional liquid-like state to a pathological gel-like or solid-like state. Here, we present a unique case to demonstrate that an unusual cascade of intermolecular charge-transfer coupled with a multitude of transient noncovalent interactions and conformational fluctuations can promote liquid phase condensation of a pH-responsive, intrinsically disordered, oligopeptide repeat domain of a melanosomal protein. At neutral cytosolic pH, the repeat domain forms highly dynamic, mesoscopic, permeable, liquid-like droplets possessing rapid internal diffusion and torsional fluctuations. These liquid condensates mature via pervasive intermolecular charge-transfer and persistent backbone interactions driving the liquid-to-solid phase transition into heterogeneous solid-like aggregates that are structurally and morphologically distinct from typical amyloids formed at mildly acidic melanosomal pH. Our findings reveal the regulatory role of the repeat domain as a specific pH-sensor that critically controls the phase transition and self-assembly processes akin to prion-like low-complexity domains modulating intracellular phase separation.
URI:	<a href="https://pubs.acs.org/doi/abs/10.1021/jacs.9b10892">https://pubs.acs.org/doi/abs/10.1021/jacs.9b10892</a> ( <a href="https://pubs.acs.org/doi/abs/10.1021/jacs.9b10892">https://pubs.acs.org/doi/abs/10.1021/jacs.9b10892</a> ) <a href="http://hdl.handle.net/123456789/1627">http://hdl.handle.net/123456789/1627</a> ( <a href="http://hdl.handle.net/123456789/1627">http://hdl.handle.net/123456789/1627</a> )
Appears in	Research Articles (/jspui/handle/123456789/9)
Collections:	

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/1627/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/1627/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/1627?mode=full>)

 (</jspui/handle/123456789/1627/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.