





Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali / Thesis & Dissertation / Master of Science / MS-19

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/5734

Title: Lattices in Euclidean space

Authors: Agrawal, Shreepad

Keywords: Lattices Euclidean

Issue Date: May-2024

Publisher: IISER Mohali

Abstract:

A lattice L is a finitely generated Z-submodule of a vector space such that it contains a basis of the vector space over Q. Given a bilinear form on L, we define a quadratic form Q(x) on the lattice. A lattice L is said to be positive lattice if Q(x) > 0 for all $x \square RV /\{0\}$. If L and M are positive lattices, we can define the tensor product L \square M which is also a positive lattice. We define the min(L) for a positive lattice to be the min{Q(x)|x} $\square L/\{0\}$ }. Then min(L $\square M$) \le min(L)min(M). The natural question is when does the equality hold. The equality holds for every M, if L is of E-type. We'll explore these special lattice and their properties. The second part of my thesis is regarding scalar extension of lattices. Let L and M be two positive lattices, F be a finite extensions of Q and R F , the ring of integers of F. Then R F \square L is called the scalar extension of L. Assume there exists an isometry σ such that σ (L) = M. Then σ is also an isometry between the scalar extensions of lattices, i.e. σ (R F \square L) = R F \square M. The interesting questions is, assume there exists an isometry between the scalar extension of lattices?

Description: under embargo period

URI: http://hdl.handle.net/123456789/5734

Appears in MS-

Collections:

Files in This Item:

File	Description	Size	Format	
Under Embargo period.odt	under embargo period	9.72 kB	OpenDocument Text	View/Open

Show full item record

alin

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.



Customized & Implemented by - Jivesna Tech