



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**

**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2653>


Title:	Quantitative characterization of new supramolecular synthons involving fluorine atoms in the crystal structures of di- and tetrafluorinated benzamides
Authors:	Yadav, H.R. (/jspui/browse?type=author&value=Yadav%2C+H.R.) Choudhury, A.R. (/jspui/browse?type=author&value=Choudhury%2C+A.R.)
Keywords:	Quantum theory of atoms in molecules (QTAIM). Intermolecular interactions Supramolecular synthons Organic fluorine
Issue Date:	2017
Publisher:	Wiley-Blackwell
Citation:	Acta Crystallographica Section B: Structural Science, Crystal Engineering and Materials, 73(5), pp. 805-819
Abstract:	Strong hydrogen bonds play a significant role in crystal packing. In particular, the involvement of interactions involving fluorine in controlling the crystal packing requires appropriate attention, especially in the presence of other strong hydrogen bonds. In the present study, a detailed quantitative assessment has been performed of the nature, energetics and topological properties derived from the electron density in model compounds based on fluorinated benzamides (a total of 46 fluorine-substituted benzamides containing multiple fluorine atoms) in the solid state. The primary motivation in the design of such molecules is to enhance the acidity of the interacting H atoms in the presence of an increasing number of F atoms on the molecular scaffold, resulting in increased propensity towards the formation of intermolecular interactions involving organic fluorine. This exercise has resulted in the identification of new and frequently occurring supramolecular synthons involving F atoms in the packing of molecules in the solid state. The energetics associated with short and directional intermolecular Csp <sup>2</sup> —H...F—Csp <sup>2</sup> interactions with significantly high electrostatic contributions is noteworthy, and the topological analysis reveals the bonding character of these ubiquitous interactions in crystal packing in addition to the presence of Csp <sup>2</sup> —F...F—Csp <sup>2</sup> contacts.
Description:	Only IISERM authors are available in the record.
URI:	<a href="https://onlinelibrary.wiley.com/iucr/doi/10.1107/S2052520617006114">https://onlinelibrary.wiley.com/iucr/doi/10.1107/S2052520617006114</a> ( <a href="https://onlinelibrary.wiley.com/iucr/doi/10.1107/S2052520617006114">https://onlinelibrary.wiley.com/iucr/doi/10.1107/S2052520617006114</a> ) <a href="http://hdl.handle.net/123456789/2653">http://hdl.handle.net/123456789/2653</a> ( <a href="http://hdl.handle.net/123456789/2653">http://hdl.handle.net/123456789/2653</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/2653/1/Need%20to%20add%20pdf.odt)		7.9 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/2653/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/2653?mode=full>)

 (</jspui/handle/123456789/2653/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.