



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/5026>


Title:	Copper(II) complexes with a benzimidazole functionalized Schiff base: Synthesis, crystal structures, and role of ancillary ions in phenoxazinone synthase activity
Authors:	Joshi, Mayank (/jspui/browse?type=author&value=Joshi%2C+Mayank) Choudhury, Angshuman Roy (/jspui/browse?type=author&value=Choudhury%2C+Angshuman+Roy)
Keywords:	Synthesis crystal structures
Issue Date:	2021
Publisher:	Wiley
Citation:	Applied Organometallic Chemistry, 35(6).
Abstract:	<p>This research study reports the synthesis, structural characterization and phenoxazinone synthase-like activity of two structurally similar copper(II) complexes developed with a benzimidazole functionalized Schiff base (L). The ligand, L, was designed and synthesized in high yield by the reaction of p-methoxy benzaldehyde with o-phenylenediamine. The reaction of L with CuCl₂ and Cu(NO₃)₂ leads to the formation of two isostructural complexes, [Cu(L)2Cl2]2 (1) and [Cu(L)2(NO3)2]2 (2). Single crystal X-ray structural study reveals that both the Cu(II) centre in 1 and 2 adopts a square planar geometry. An attempt has also been made to understand the role of coordinated co-ligands on the catalytic oxidation of 2-aminophenol (2-AP) to 2-amino-3H-phenoxazine-3-one (2-APX) in methanol. The presence of coordinated nitrate to Cu(II) ions imparts a more labile character to complex 2, and the catalytic efficiency (kcat/KM) for complex 2 (1.50 × 10⁷) was determined almost double compared with that of complex 1 (8.78 × 10⁶). Electro-chemical and electrospray ionization mass spectrometry studies of 1 and 2 with 2-AP suggests that the square planar geometries of the Cu(II) centres remain the driving force to develop enzyme-substrate adducts and excellent catalytic performance of the complexes. Electrochemical and EPR spectral analysis of the reaction mixture confirm the presence of active 2-AP-/2-AP+- redox species in the course of catalytic oxidation and suggest the radical driven oxidative coupling of 2-AP in an aerobic environment. Temperature-dependent kinetic measurements were carried out to evaluate the activation parameters (E_a, ΔH‡, ΔS‡), which favours the higher rate of catalytic oxidation of 2-AP for complex 2 than complex 1.</p>
Description:	Only IISER Mohali authors are available in the record.
URI:	https://onlinelibrary.wiley.com/doi/10.1002/aoc.6211 (https://onlinelibrary.wiley.com/doi/10.1002/aoc.6211) http://hdl.handle.net/123456789/5026 (http://hdl.handle.net/123456789/5026)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need To Add...Full Text_PDF (/jspui/bitstream/123456789/5026/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF)		15.36 kB	Unknown

View/Open (/jspui/t

Show full item record (</jspui/handle/123456789/5026?mode=full>)

 (</jspui/handle/123456789/5026/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.