



# Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali / Thesis & Dissertation / Master of Science / MS-19

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/5707>

Title:	The Regulation of NMDA Receptors at GABAergic Interneurons During Postnatal Development
Authors:	<a href="#">Paul, Soumyadev</a>
Keywords:	D-serine Glycine NMDA
Issue Date:	May-2024
Publisher:	IISER Mohali
Abstract:	Brain development is marked by periods of enhanced brain plasticity, known as critical periods. The termination of critical periods requires the maturation of parvalbumin-positive interneurons (PV + INs). Several studies have linked the disruption of N-methyl D-aspartate receptors (NMDARs) in PV + INs during postnatal development to the pathophysiology of schizophrenia and autism spectrum disorders. Canonical NMDA receptors are hetero-tetramers with two GluN1 and two GluN2 subunits, which together form an ion channel. GluN2 subunits can be of four classes- GluN2A/B/C/D. NMDAR activation requires the simultaneous binding of the excitatory neurotransmitter glutamate, and a co-agonist, which can be glycine or D-serine. Recent studies by our lab have found that in adult mice, D-serine but not glycine is critical for maintaining the activity of NMDARs at PV INs in the late adolescent-young adult prefrontal cortex (PFC), and that loss of D-serine functions leads to the synaptic deficits observed in neuropsychiatric disorders such as schizophrenia. However, it is not known if the identity of the NMDAR co-agonist in these cells is developmentally regulated. Through this study, I aim to test the functions of D-serine and glycine as NMDAR co-agonists throughout postnatal development using the transgenic PV -tdTomato mice where PV INs are readily identifiable. Using selective enzymatic scavengers to block the function of either D-serine or glycine, I reveal that D-serine but not glycine gates NMDARs at the PFC PV INs at juvenile synapses (11-17 day old mice). Strikingly, I also show that bath application of D-serine inhibits NMDA-EPSCs at PV+ in neonates (11-17 days-old) while it does increase the synaptic responses in mature synapses (45-70 days old). Finally, I test the hypothesis for a change in the composition of NMDARs subunits in PFC PV INs during development. Overall, this study helps in understanding the relative contribution of D-serine and glycine in the regulation of specific NMDARs at PV INs during postnatal development.
Description:	Under Embargo Period
URI:	<a href="http://hdl.handle.net/123456789/5707">http://hdl.handle.net/123456789/5707</a>
Appears in Collections:	<a href="#">MS-19</a>

## Files in This Item:

File	Description	Size	Format	
<a href="#">embargo period.pdf</a>		6.04 kB	Adobe PDF	<a href="#">View/Open</a>

Show full item record



Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.