



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4813>


Title:	Protocol for investigating the biogenesis of SARS-CoV-2 S pseudoviruses in HEK293T cells transduced to express the virus-specific intrabodies
Authors:	Dahiya, Surbhi (/jspui/browse?type=author&value=Dahiya%2C+Surbhi) Singh, Sudhakar (/jspui/browse?type=author&value=Singh%2C+Sudhakar) Sehrawat, Sharvan (/jspui/browse?type=author&value=Sehrawat%2C+Sharvan)
Keywords:	SARS-CoV-2 S pseudoviruses HEK293T cells virus-specific intrabodies
Issue Date:	2022
Publisher:	Elsevier
Citation:	STAR Protocols, 4(1), 1-14.
Abstract:	<p>This protocol provides detailed methodology used in the published article; Robust anti-SARS-CoV-2 single domain antibodies cross neutralize multiple viruses.1 The protocol describes a strategy for the intracellular expression of sdAbs targeting the polybasic cleavage site (anti-CSP sdAb) of the viral entry mediator, SARS-CoV-2 S protein. We also analyze the intracellular fate of the protein in the anti-CSP intrabody (IB)-expressing cells. For generating an anti-CSP IB construct, the sequence of the selected sdAb specific to peptide (NSPRRAR/SVAS) which encompasses the polybasic cleavage site (CSP) of SARS-CoV-2 S protein was cloned into a pLenti-GFP vector (a lentivirus-based mammalian expression vector) downstream to CMV promoter.2 The cloning scheme used is: 50 LTR-CMV promoter-VHH - c-Myc-tag - e-GFP -30 - LTR. The anti-CSP sdAb sequence was amplified using the forward primer, VHH-FR1-pLenti (containing XbaI restriction site), and reverse primer, FR4-c-Myc pLenti, that encoded c-Myc tag at 30 end of VHH with a BamHI site) (Figure 1). To generate the lentivirus-based pseudoviruses expressing surface SARS-CoV-2 S protein, HEK293T cells were transfected with plasmids such as a reporter construct, pCMVR8.74, SARS-CoV-2 S construct, tat and rev as well as pMD2.G.3 The plasmids required for producing LV(CoV-2 S) pseudoviruses were prepared and the purity was confirmed before beginning with the experiments.</p>
Description:	Only IISERM authors are available in the record
URI:	https://doi.org/10.1016/j.xpro.2022.101977 (https://doi.org/10.1016/j.xpro.2022.101977) http://hdl.handle.net/123456789/4813 (http://hdl.handle.net/123456789/4813)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need To Add...Full Text_PDF (/jspui/bitstream/123456789/4813/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF)		15.36 kB	Unknown

[View/Open \(/jspui/t](#)

Show full item record (</jspui/handle/123456789/4813?mode=full>)

 (</jspui/handle/123456789/4813/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.