



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4400>


Title:	Tyrosine in the hinge region of the pore-forming motif regulates oligomeric β -barrel pore formation by <i>Vibrio cholerae</i> cytolysin
Authors:	Mondal, Anish Kumar (/jspui/browse?type=author&value=Mondal%2C+Anish+Kumar) Verma, Paras (/jspui/browse?type=author&value=Verma%2C+Paras) Sengupta, Nayanika (/jspui/browse?type=author&value=Sengupta%2C+Nayanika) Dutta, Somnath (/jspui/browse?type=author&value=Dutta%2C+Somnath) Pandit, Shashi Bhushan (/jspui/browse?type=author&value=Pandit%2C+Shashi+Bhushan) Chattopadhyay, Kausik (/jspui/browse?type=author&value=Chattopadhyay%2C+Kausik)
Keywords:	Tyrosine hinge region pore-forming oligomeric β -barrel
Issue Date:	2021
Publisher:	Wiley
Citation:	Molecular Microbiology, 115(4), 508-525.
Abstract:	β -barrel pore-forming toxins perforate cell membranes by forming oligomeric β -barrel pores. The most crucial step is the membrane-insertion of the pore-forming motifs that create the transmembrane β -barrel scaffold. Molecular mechanism that regulates structural reorganization of these pore-forming motifs during β -barrel pore-formation still remains elusive. Using <i>Vibrio cholerae</i> cytolysin as an archetypical example of the β -barrel pore-forming toxin, we show that a key tyrosine residue (Y321) in the hinge region of the pore-forming motif plays crucial role in this process. Mutation of Y321 abrogates oligomerization of the membrane-bound toxin protomers, and blocks subsequent steps of pore-formation. Our study suggests that the presence of Y321 in the hinge region of the pore-forming motif is crucial for the toxin molecule to sense membrane-binding, and to trigger essential structural rearrangements required for the subsequent oligomerization and pore-formation process. Such a regulatory mechanism of pore-formation by <i>V. cholerae</i> cytolysin has not been documented earlier in the structurally related β -barrel pore-forming toxins.
Description:	Only IISER Mohali authors are available in the record.
URI:	https://doi.org/10.1111/mmi.14631 (https://doi.org/10.1111/mmi.14631) http://hdl.handle.net/123456789/4400 (http://hdl.handle.net/123456789/4400)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format

Need To Add...Full Text_PDF..pdf (/jspui/bitstream/123456789/4400/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF..pdf)	Only IISER Mohali authors are available in the record.	15.36 kB	Adobe PDF	View/Open (/jspu
--	--	-------------	--------------	----------------------------------

[Show full item record \(/jspui/handle/123456789/4400?mode=full\)](#)

 [\(/jspui/handle/123456789/4400/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.