



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4826>


Title:	Designed pincer ligand supported Co(ii)-based catalysts for dehydrogenative activation of alcohols: Studies on N-alkylation of amines, α -alkylation of ketones and synthesis of quinolines
Authors:	Singh, Anshu (/jspui/browse?type=author&value=Singh%2C+Anshu) Maji, Ankur (/jspui/browse?type=author&value=Maji%2C+Ankur) Joshi, Mayank (/jspui/browse?type=author&value=Joshi%2C+Mayank) Choudhury, Angshuman R. (/jspui/browse?type=author&value=Choudhury%2C+Angshuman+R.) Ghosh, Kaushik (/jspui/browse?type=author&value=Ghosh%2C+Kaushik)
Keywords:	Co(ii)-based catalysts N-alkylation
Issue Date:	2021
Publisher:	Publishing
Citation:	Dalton Transactions. 50, 8567-8587.
Abstract:	Base-metal catalysts Co1, Co2 and Co3 were synthesized from designed pincer ligands L1, L2 and L3 having NNN donor atoms respectively. Co1, Co2 and Co3 were characterized by IR, UV-Vis. and ESI-MS spectroscopic studies. Single crystal X-ray diffraction studies were investigated to authenticate the molecular structures of Co1 and Co3. Catalysts Co1, Co2 and Co3 were utilized to study the dehydrogenative activation of alcohols for N-alkylation of amines, α -alkylation of ketones and synthesis of quinolines. Under optimized reaction conditions, a broad range of substrates including alcohols, anilines and ketones were exploited. A series of control experiments for N-alkylation of amines, α -alkylation of ketones and synthesis of quinolines were examined to understand the reaction pathway. ESI-MS spectral studies were investigated to characterize cobalt-alkoxide and cobalt-hydride intermediates. Reduction of styrene by evolved hydrogen gas during the reaction was investigated to authenticate the dehydrogenative nature of the catalysts. Probable reaction pathways were proposed for N-alkylation of amines, α -alkylation of ketones and synthesis of quinolines on the basis of control experiments and detection of reaction intermediates.
Description:	Only IISERM authors are available in the record
URI:	https://pubs.rsc.org/en/content/articlelanding/2021/DT/D0DT03748F (https://pubs.rsc.org/en/content/articlelanding/2021/DT/D0DT03748F) http://hdl.handle.net/123456789/4826 (http://hdl.handle.net/123456789/4826)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need To Add...Full Text_PDF (/jspui/bitstream/123456789/4826/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF)		15.36 kB	Unknown

[View/Open \(/jspui/t](#)

Show full item record (</jspui/handle/123456789/4826?mode=full>)

 (</jspui/handle/123456789/4826/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.