



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**  
**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**  
**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4471>


Title:	Expanding the Utility of Inexpensive Pyridine-N-oxide Directing Group for the Site-selective sp <sup>2</sup> /sp <sup>3</sup> γ-C-H and sp <sup>2</sup> δ-C-H Functionalization of Carboxamides
Authors:	Tomar, Radha (/jspui/browse?type=author&value=Tomar%2C+Radha) Kumar, Amit (/jspui/browse?type=author&value=Kumar%2C+Amit) Dalal, Arup (/jspui/browse?type=author&value=Dalal%2C+Arup) Bhattacharya, Debabrata (/jspui/browse?type=author&value=Bhattacharya%2C+Debabrata) Singh, Prabhakar (/jspui/browse?type=author&value=Singh%2C+Prabhakar) Babu, Srinivasarao Arulananda (/jspui/browse?type=author&value=Babu%2C+Srinivasarao+Arulananda)
Keywords:	Pyridine-N-oxide Carboxamides
Issue Date:	2022
Publisher:	Wiley
Citation:	Asian Journal of Organic Chemistry, 11(9), 2200311
Abstract:	We have shown our efforts toward expanding the utility of the relatively inexpensive pyridine-N-oxide directing group in the Pd(II)-catalyzed site-selective γ-C(sp <sup>2</sup> )-H, γ-C(sp <sup>3</sup> )-H and δ-C(sp <sup>2</sup> )-H functionalization. The functionalization β-C-H bonds using bidentate directing group (DG) pyridine-N-oxide which operates through the N,O-coordination mode has been well documented in the literature. However, there exist rare reports dealing with the functionalization of remote sp <sup>2</sup> /sp <sup>3</sup> γ- and δ-C-H bonds of carboxamides assisted by the bidentate directing groups operating via the N,O-coordination. In this paper, the scope of pyridine-N-oxide DG was examined for accomplishing the site-selective (mono) γ-C(sp <sup>2</sup> )-H arylation in substrates containing competitive C(sp <sup>3</sup> )-H and C(sp <sup>2</sup> )-H bonds. The investigation has enabled to assemble a library of pyridine-N-oxide-based biarylacetamides, heteroaryl-based biaryl carboxamides, tricyclic quinolones, arylheteroarylmethanes, biaryl-based aliphatic carboxamides and mono (ortho) arylated phenylglycine derivatives. In general, biaryl derivatives and in particular, arylacetamide, arylacetic acid derivatives and pyridine-N-oxide (2-aminopyridyl) motifs are medicinally relevant classes of compounds. This work enabled the assembling of a library of the above-mentioned types of compounds through the pyridine-N-oxide directing group-aided site-selective sp <sup>2</sup> /sp <sup>3</sup> γ-C-H and sp <sup>2</sup> δ-C-H functionalization of carboxamides.
Description:	Only IISERM authors are available in the record
URI:	<a href="https://doi.org/10.1002/ajoc.202200311">https://doi.org/10.1002/ajoc.202200311</a> ( <a href="https://doi.org/10.1002/ajoc.202200311">https://doi.org/10.1002/ajoc.202200311</a> ) <a href="http://hdl.handle.net/123456789/4471">http://hdl.handle.net/123456789/4471</a> ( <a href="http://hdl.handle.net/123456789/4471">http://hdl.handle.net/123456789/4471</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need To Add...Full Text_PDF. (/jspui/bitstream/123456789/4471/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF.)		15.36 kB	Unknown

View/Open (/jspui/

Show full item record (</jspui/handle/123456789/4471?mode=full>)

 (</jspui/handle/123456789/4471/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.