

Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

- / Publications of IISER Mohali (/jspui/handle/123456789/4)
- / Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/1653

Title: Synthesis and spectroscopic characterization of a photo-stable tetrazinc(II)–Schiff base cluster:

A rare case of ligand centric phenoxazinone synthase activity

Authors: Choudhury, A.R. (/jspui/browse?type=author&value=Choudhury%2C+A.R.)

Joshi, Mayank (/jspui/browse?type=author&value=Joshi%2C+Mayank)

Keywords: Zinc(II)

Schiff base X-ray structure Bio-mimicking study

Phenoxazinone synthase activity

Issue Date: 2018

Publisher: ScienceDirect.

Citation: Polyhedron, 156, pp. 223-230

Abstract:

Herein, the synthesis, structural characterization and catalytic activity of a novel tetranuclear zinc(II)—Schiff base complex, [Zn4(L)2(μ 3-OCH3)2(CH3OH)2]·2CH3OH (1), [L = N,N'-bis(3methoxysalicylidene)-1,3-diamino-2-propanol] was presented. Single crystal X-ray diffraction structural analysis revealed that the tetra-zinc(II) cluster crystallized in a monoclinic system with P21/c space group. Interestingly, three different molecular bridges (methoxido-, alkoxido- and phenoxido-) simultaneously co-existed in assembling tetra-zinc(II) core, which was a very rare observation. To the best of our knowledge, this compound would be the first compound where a diverse coordination aspect was covered by a single solvent as terminal coordinator (CH3OH), bridging (µ3-CH3OH) and solvent for crystallization in the existing scientific literature. The compound showed good photo-stability and excellent luminescence property with higher lifetime at transition state in ethanol. This zinc(II) complex revealed crucial role as an effective catalytic system towards oxidation of 2-aminophenol (2-AP) in ethanol. Additionally, the tetra-zinc(II) complex displayed potential phenoxazinone synthase like activity with momentous turn over number, kcat $(h-1) = 6.19 \times 102$ in ethanol under aerobic condition. ESI-MS and EPR spectral analysis of the reaction mixture between Zn(II) complex and 2-AP recommended that the course of catalysis proceeded through substrate-catalyst adduct formation and authenticated the radical mechanistic pathway in favor of oxidative coupling product. This tetranuclear zinc(II)-Schiff base complex would be considered as the first example that catalyzed the oxidative coupling of 2aminophenol to aminophenoxazino compound under usual aerobic condition. As complementary, detailed quantum chemical computations, performed with density functional theory (DFT) were well corroborated with the experimental results. This was the first and rarest example where a tetrazinc(II)-Schiff base cluster exhibited catalytic oxidation of 2-AP through ligand centered radical activity.

Description: Only IISERM authors are available in the record.

URI: https://www.sciencedirect.com/science/article/pii/S0277538718305965

(https://www.sciencedirect.com/science/article/pii/S0277538718305965)

http://hdl.handle.net/123456789/1653 (http://hdl.handle.net/123456789/1653)

Appears in Collections:

Research Articles (/jspui/handle/123456789/9)

Files in This Item:				
File	Description	Size	Format	
Need to add pdf.odt (/jspui/bitstream/123456789/1653/1/Need%20to%20add%20pdf.odt)		8.04 kB	OpenDocument Text	View/Open (/jspui/bitstream/12345

Show full item record (/jspui/handle/123456789/1653?mode=full)

. (/jspui/handle/123456789/1653/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.