



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4706>


Title:	Thermal transport of the frustrated spin-chain mineral linarite: Magnetic heat transport and strong spin-phonon scattering
Authors:	Sakrikar, Piyush (/jspui/browse?type=author&value=Sakrikar%2C+Piyush)
Keywords:	Frustrated magnetism Magnons Lattice thermal conductivity
Issue Date:	2021
Publisher:	American Physical Society
Citation:	Physical Review B, 104(23).
Abstract:	The mineral linarite $[\text{PbCuSO}_4(\text{OH})_2]$ forms a monoclinic structure where a sequence of $\text{Cu}(\text{OH})_2$ units forms a spin-1/2 chain. Competing ferromagnetic nearest-neighbor (J_1) and antiferromagnetic next-nearest-neighbor interactions (J_2) in this quasi-one-dimensional spin structure imply magnetic frustration and lead to magnetic ordering below $T_N = 2.8$ K in a multiferroic elliptical spin-spiral ground state. Upon the application of a magnetic field along the spin-chain direction, distinct magnetically ordered phases can be induced. We studied the thermal conductivity κ in this material across the magnetic phase diagram as well as in the paramagnetic regime in the temperature ranges 0.07–1 K and 9–300 K. We found that in linarite the heat is carried mainly by phonons but shows a peculiar nonmonotonic behavior in field. In particular, κ is highly suppressed at the magnetic phase boundaries, indicative of strong scattering of the phonons off critical magnetic fluctuations. Even at temperatures far above the magnetically ordered phases, the phononic thermal conductivity is reduced due to scattering off magnetic fluctuations. The mean free path due to spin-phonon scattering ($l_{\text{spin-phonon}}$) was determined as a function of temperature. A power law behavior was observed mainly above 500 mK indicating the thermal activation of spin fluctuations. In the critical regime close to the saturation field, $l_{\text{spin-phonon}}$ shows a $1/T$ dependence. Furthermore, a magnon thermal transport channel was verified in the helical magnetic phase. We estimate a magnon mean free path which corresponds to about 1000 lattice spacings.
Description:	Only IISERM authors are available in the record
URI:	https://doi.org/10.1103/PhysRevB.104.235129 (https://doi.org/10.1103/PhysRevB.104.235129) http://hdl.handle.net/123456789/4706 (http://hdl.handle.net/123456789/4706)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need To Add...Full Text_PDF (/jspui/bitstream/123456789/4706/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF)		15.36 kB	Unknown

[View/Open \(/jspui/t\)](#)

Show full item record (</jspui/handle/123456789/4706?mode=full>)

 (</jspui/handle/123456789/4706/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.