



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1931>


Title:	Coordination polymers of CdII and PbII derived from bipyridine–glycoluril: influence of metal-ion size and counter-ions
Authors:	Markad, D. (/jspui/browse?type=author&value=Markad%2C+D.)
Keywords:	Bipyridine-glycoluril Coordination polymer Crystal structure Hydrogen bonding
Issue Date:	2019
Publisher:	Acta Crystallographica Section C
Citation:	Acta Crystallographica Section C: Structural Chemistry,75, pp. 1084-1090.
Abstract:	Two new one-dimensional (1D) coordination polymers (CPs), namely catena-poly[[[aqua-cadmium(II)]-bis(μ -4b,5,7,7a-tetrahydro-4b,7a-epiminomethanoimino-6H-imidazo[4,5-f][1,10]phenanthroline-6,13-dione)] bis(perchlorate) dihydrate], {[Cd(C ₁₄ H ₁₀ N ₆ O ₂) ₂ (H ₂ O)](ClO ₄) ₂ ·2H ₂ O} _n or {[Cd(BPG) ₂ (H ₂ O)](ClO ₄) ₂ ·2H ₂ O} _n , 1, and catena-poly[[lead(II)-bis(μ -4b,5,7,7a-tetrahydro-4b,7a-epiminomethanoimino-6H-imidazo[4,5-f][1,10]phenanthroline-6,13-dione)] bis(perchlorate) dihydrate], {[Pb(C ₁₄ H ₁₀ N ₆ O ₂) ₂](ClO ₄) ₂ ·2H ₂ O} _n or {[Pb(BPG) ₂](ClO ₄) ₂ ·2H ₂ O} _n , 2, have been synthesized using bipyridine–glycoluril (BPG; systematic name: 4b,5,7,7a-tetrahydro-4b,7a-epiminomethanoimino-6H-imidazo[4,5-f][1,10]phenanthroline-6,13-dione), a urea-fused tecton, in a mixed-solvent system. The CdII ion in 1 is heptacoordinated and the PbII ion in 2 is hexacoordinated, with the CdII ion adopting a pentagonal bipyramidal geometry and the PbII ion adopting a distorted octahedral geometry. Both CPs form infinite linear chain structures which are hydrogen bonded to each other leading to the formation of three-dimensional supramolecular network structures. Topological analysis of CPs 1 and 2 reveals that the structures exhibit 1D chain-like arrangements in an AB–AB sequence and shows platonic uniform 2-connected uninodal topologies. Furthermore, a comparative analysis of a series of structures based on the BPG ligand indicates that the size of the metal ion and the types of counter-ions used have a great influence on the resulting frameworks and properties.
Description:	Only IISERM authors are available in the record.
URI:	http://hdl.handle.net/123456789/1931 (http://hdl.handle.net/123456789/1931) https://onlinelibrary.wiley.com/iucr/doi/10.1107/S2053229619008325 (https://onlinelibrary.wiley.com/iucr/doi/10.1107/S2053229619008325)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/1931/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/1931/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/1931?mode=full>)

 (</jspui/handle/123456789/1931/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.