



## Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Thesis & Dissertation (/jspui/handle/123456789/1)**

**/ Doctor of Philosophy (PhD) (/jspui/handle/123456789/268)**

**/ PhD-2013 (/jspui/handle/123456789/269)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1221>

**Title:** Effect of Chemical Substitutions on the Quantum Spin Liquid Candidates  $\text{Na}_4\text{Ir}_3\text{O}_8$  and  $\text{Ca}_{10}\text{Cr}_7\text{O}_{28}$

**Authors:** Balodhi, A. (/jspui/browse?type=author&value=Balodhi%2C+A.)

**Keywords:** Physics  
Quantum Spin  
X-ray diffraction  
Magnetic measurement  
Resistivity

**Issue Date:** 1-Oct-2019

**Publisher:** IISERM

**Abstract:** Quantum spin liquids (QSLs) are exotic states of matter, in which magnetic frustration and strong quantum fluctuations destroy long-range magnetic order. Highly frustrated lattices with antiferromagnetic exchange interactions and low-spin value open up possibilities in the exploration and designing of new QSL candidates. QSLs in three-dimensions are very rare, and only a couple of candidate materials exist. In this thesis, we investigate bond-disordered quantum spin-liquid state in three dimensional magnetic insulators  $\text{Na}_4\text{Ir}_3\text{O}_8$  and  $\text{Ca}_{10}\text{Cr}_7\text{O}_{28}$ . The central theme of thesis is to explore how these QSL's respond to external perturbations like pressure, magnetic field, and chemical substitutions. In this thesis, I have synthesized two QSL candidates the hyperkagome  $\text{Na}_4\text{Ir}_3\text{O}_8$  and kagome bilayer  $\text{Ca}_{10}\text{Cr}_7\text{O}_{28}$  and studied their electrical transport, magnetic, and thermal properties. Our work provides several new results: (i) the strongly frustrated Mott insulating state in  $\text{Na}_4\text{Ir}_3\text{O}_8$  is quite robust against large removal of Na from the lattice, (ii) evidence through magnetic (Ru) and nonmagnetic (Ti) impurity doping in a magnetic sublattice, of fragile magnetic order and importance of nearestneighbour interactions and spin-orbit coupling in deciding the magnetic ground state in  $\text{Na}_4\text{Ir}_3\text{O}_8$ , (iii) high-pressure magnetic susceptibility measurements reveal the QSL at ambient pressure is quite robust and may not depend on a delicate balance between any specific values of competing exchange interactions in  $\text{Ca}_{10}\text{Cr}_7\text{O}_{28}$ , and (iv) the first experimental realization of a perfect nonmagnetic analog  $\text{Ca}_{10}\text{V}_7\text{O}_{27.5}$  of  $\text{Ca}_{10}\text{Cr}_7\text{O}_{28}$ , which enables an accurate exclusion of the lattice heat capacities.

**URI:** IISERM (IISERM)  
<http://hdl.handle.net/123456789/1221> (<http://hdl.handle.net/123456789/1221>)

**Appears in** PhD-2013 (/jspui/handle/123456789/269)  
**Collections:**

Files in This Item:

File	Description	Size	Format	
PH13009.pdf (/jspui/bitstream/123456789/1221/4/PH13009.pdf)		36.87 MB	Adobe PDF	<a href="#">View/Open (/jspui/bitstream/123456789/1221/4,</a>

[Show full item record \(/jspui/handle/123456789/1221?mode=full\)](#)

[📊 \(/jspui/handle/123456789/1221/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

Admin Tools

[Edit...](#)

[Export Item](#)

[Export \(migrate\) Item](#)

[Export metadata](#)