



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2504>


Title:	Characterization of Salt-Induced Oligomerization of Human β 2-Microglobulin at Low pH
Authors:	Narang, D. (/jspui/browse?type=author&value=Narang%2C+D.) Singh, Anubhuti (/jspui/browse?type=author&value=Singh%2C+Anubhuti) Hemaswathi, M. (/jspui/browse?type=author&value=Hemaswathi%2C+M.) Mukhopadhyay, S. (/jspui/browse?type=author&value=Mukhopadhyay%2C+S.)
Keywords:	Amyloidosis Misfolding Aggregation Amyloid
Issue Date:	2016
Publisher:	American Chemical Society
Citation:	Journal of Physical Chemistry B 120(32), pp.7815-7823.
Abstract:	Misfolding and amyloid aggregation of human β 2-microglobulin (β 2m) have been linked to dialysis-related amyloidosis. Previous studies have shown that in the presence of different salt concentrations and at pH 2.5, β 2m assembles into aggregates with distinct morphologies. However, the structural and mechanistic details of the aggregation of β 2m, giving rise to different morphologies, are poorly understood. In this work, we have extensively characterized the salt-induced oligomers of the acid-unfolded state of β 2m using an array of biophysical tools including steady-state and time-resolved fluorescence, circular dichroism, dynamic light scattering, and atomic force microscopy imaging. Fluorescence studies using the oligomer-sensitive molecular rotor, 4-(dicyanovinyl)-julolidine, in conjunction with the light scattering and cross-linking assay indicated that at low salt (NaCl) concentrations β 2m exists as a disordered monomer, capable of transforming into ordered amyloid. In the presence of higher concentrations of salt, β 2m aggregates into a larger oligomeric species that does not appear to transform into amyloid fibrils. Site-specific fluorescence experiments using single Trp variants of β 2m revealed that the middle region of the protein is incorporated into these oligomers, whereas the C-terminal segment is highly exposed to bulk water. Additionally, stopped-flow kinetic experiments indicated that the formation of hydrophobic core and oligomerization occur concomitantly. Our results revealed the distinct pathways by which β 2m assembles into oligomers and fibrils.
URI:	https://pubs.acs.org/doi/10.1021/acs.jpcb.6b05619 (https://pubs.acs.org/doi/10.1021/acs.jpcb.6b05619) http://hdl.handle.net/123456789/2504 (http://hdl.handle.net/123456789/2504)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/2504/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/2504/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/2504?mode=full>)

 (</jspui/handle/123456789/2504/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.