



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1998>

Title:	Label-Free Imaging of Fibronectin Adsorption at Poly-(l-lysine)-Decorated Liquid Crystal Droplets
Authors:	Verma, I. (/jspui/browse?type=author&value=Verma%2C+I.) Pani, I. (/jspui/browse?type=author&value=Pani%2C+I.) Sharma, Diksha (/jspui/browse?type=author&value=Sharma%2C+Diksha) Maity, Shiny (/jspui/browse?type=author&value=Maity%2C+Shiny) Pal, S.K. (/jspui/browse?type=author&value=Pal%2C+S.K.)
Keywords:	Fibronectin Experimental system Interfacial systems
Issue Date:	2019
Publisher:	American Chemical Society
Citation:	Journal of Physical Chemistry C, 123(22),pp. 13642-13650.
Abstract:	Interactions and subsequent structural organization of fibronectin (FibN) (major component of an extracellular matrix) molecules in contact with chemically modified surfaces form an essential basis to the understanding of the regulation of matrix assembly in pathological states and to the development of new biomedical devices. Here, using poly-l-lysine (PLL)-coated liquid crystal (LC) droplets, we present a simple but fruitful advance of an experimental system that can report label-free imaging of adsorption of FibN at the LC–aqueous interface. We observed that the interfacial intermolecular interactions of anionic FibN molecules and cationic PLL residues is accompanied by a director configuration transition of the LC droplets from radial to bipolar/preradial, giving a distinct optical output. Specifically, it is found that the ordering transitions of LC can be effectively tuned by blocking the anionic sites of FibN with divalent cations (Ca ²⁺). Combination of fluorescence measurements, circular dichroism, and atomic force microscopy further revealed that PLL induces significant conformational changes in FibN at the aqueous–LC interface. This report provides a simple method based on LC droplets to understand polymer–protein interactions at membrane interfaces which would have potential applications in biomedical and interfacial systems.
URI:	https://pubs.acs.org/doi/10.1021/acs.jpcc.9b01934 (https://pubs.acs.org/doi/10.1021/acs.jpcc.9b01934) http://hdl.handle.net/123456789/1998 (http://hdl.handle.net/123456789/1998)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/1998/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/1998/1/Need%20to%20add%20pdf.odt\)](#)

[Show full item record \(/jspui/handle/123456789/1998?mode=full\)](#)

[Statistics \(/jspui/handle/123456789/1998/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.