



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)


Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/160>

Title:	First report on N,N'-diisoalkylisonicotinamide 1D coordination network containing linear trinuclear [CO ₃ L ₄ Cl ₆] units with mixed Coll(Td)-Coll(Oh)-Coll(Td) geometries: Structure and magnetic properties
Authors:	Kapoor, Ramesh (/jspui/browse?type=author&value=Kapoor%2C+Ramesh)
Keywords:	Carbonyl oxygen atoms Coordination networks Coordination Polymers
Issue Date:	2010
Publisher:	The Royal Society of Chemistry.
Citation:	Dalton Transactions, 39 (34), pp. 7951-7959.
Abstract:	Reaction of anhydrous CoCl ₂ with N,N'- diisopropylisonicotinamide (L) has yielded a coordination polymer containing linear trinuclear [Co ₃ L ₄ Cl ₆] units with a rare, mixed Co(Td)-Co(Oh)-Co(Td) assembly (compound 1). The central Coll ion, of each trinuclear entity, exhibits a distorted octahedral geometry, with two ligand molecules coordinating through their carbonyl oxygen atoms along with two bridging Cl ⁻ ions and two pyridine N atoms from the neighboring molecules. Also, in each unit, two outer Coll ions display distorted tetrahedral geometry, coordinating to one ligand molecule through the pyridine N atom and to three Cl ⁻ ions (one of them bridged to the central Coll and the two acting as a terminal ligands). The magnetic properties of this compound were investigated in the temperature range of 2.0 to 300.0 K. Owing to the complexity of the system and the weak interactions among trinuclear aggregates, the magnetic response has been analyzed using a model which considers these units as isolated systems. In addition, magnetic data has been examined in two separated blocks, above and below 50 K, applying programs VPMAG FORTRAN and MAGPACK-fit, respectively. This way, only the most significant effects at each interval of temperature were considered: spin-orbit coupling of the Co(Oh), at high temperatures and zero-field splitting parameters of the Co(Td) at the low. Spin-spin magnetic interaction has been taken into account for the whole range of temperatures. As a result, the analysis of the magnetic data shows that, within every trinuclear unit, the central position matches well with a high-spin Coll (S = 3/2) and also reveals weak ferromagnetic interactions between the Co(Oh) and the two terminal Co(Td) ions (J = +0.34 cm ⁻¹).
Description:	Only IISERM authors are available in the record.
URI:	http://www.ncbi.nlm.nih.gov/pubmed/20657942 (http://www.ncbi.nlm.nih.gov/pubmed/20657942) http://pubs.rsc.org/en/Content/ArticleLanding/2010/DT/c0dt00245c (http://pubs.rsc.org/en/Content/ArticleLanding/2010/DT/c0dt00245c)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format	
Need to add pdf.odt (/jspui/bitstream/123456789/160/3/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text	View/Open (/jspui/bitstream/123456789/160/3/Need%20to%20add%20pdf.odt)

Show full item record (</jspui/handle/123456789/160?mode=full>)

 (</jspui/handle/123456789/160/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.