



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2122>

| | |
|-------------------------|---|
| Title: | Crystal Engineering and Magnetostructural Properties of Newly Designed Azide/Acetate-Bridged Mn ₁₂ Coordination Polymers |
| Authors: | Ali, Anzar (/jspui/browse?type=author&value=Ali%2C+Anzar) |
| Keywords: | Crystal Engineering Coordination Polymers |
| Issue Date: | 2019 |
| Publisher: | ACS Publications |
| Citation: | Crystal Growth and Design, 19(4),pp. 2366-2379. |
| Abstract: | <p>Crystal engineering of the coordination polymers where polynuclear clusters are building blocks constitutes an emerging class of chemistry. The fine-tuning of the structural motifs leads to interesting and varying magnetic properties. Owing to such properties, two rare μ_6-oxo centered mixed-valent, azide or acetate-bridged coordination polymers viz, $[\{Mn^{II}2Mn^{III}10Na_2(\mu_6-O)_2(N_3)_{10}(NO_3)(H_2O)_4(thme)_8\} \cdot 3(Et_3NH)]_n$ (1) and $[\{Mn^{II}3Mn^{III}9Na_7(\mu_2-O)_2(\mu_6-O)_2(O)_5(CH_3O)(CH_3CO_2)_{11}(thmp)_8\} \cdot 4(O)]_n$ (2), with retention of a Mn₁₂ metallic core in both polymers are obtained using tripodal polyalcohol 1,1,1-tris (hydroxymethyl)ethane (H₃thme) and 1,1,1-tris (hydroxymethyl)propane (H₃thmp) ligands, respectively. X-ray analysis shows that 1 is a one-dimensional coordination polymer where Mn₁₂ units are propagating by bridging azide function. 1 shows the underlying net of 2,2,3C₆ topological type. 2 forms a cyclic ring as a result of repeating Mn₁₂ zigzag chains bridging by sodium and H₂O. The topology of 2 results in a 31-nodal underlying 3,3,3,3,4,4,4,4,4,4,4,4,4,4,5,5,5,5,5,5,6,6,6,6,6,6,6,6,7,7-c net with a point symbol of the net of {3.4.5}2{3.47.52}2{32.410.52.6}{32.410.53}4{32.43.5.203.21}{32.43.5}6{32.44}2{32.46.52.63.7.8}{32.46.52}{34.44.52}{34.46.54.64.72.8}2{412.52.6}2{42.6}2{43}{45.5}3{45.6}{46.63.8}. In 1, a magnetic study ascertains the presence of antiferromagnetic interaction and shows single molecule magnet-like behavior with an energy barrier of 75.5 K. However, 2 exhibited strong antiferromagnetic interaction in dc studies. The super-paramagnetic-like slow relaxation of its magnetization was not observed for 2 in out-of-phase ac magnetic susceptibility due to the absence of a large enough energy barrier. Magnetization versus an applied dc field exhibited a hysteresis loop at 2 K with a coercivity of 1069.10 Oe and remanent magnetization of 0.374 μB in 1, while 2 has no coercivity in the hysteresis loop even at the lowest temperature (2.0 K), and no saturation was observed up to 7.0 T field supporting antiferromagnetic interactions present in the polymer.</p> |
| Description: | Only IISERM authors are available in the record. |
| URI: | https://pubs.acs.org/doi/10.1021/acs.cgd.9b00058 (https://pubs.acs.org/doi/10.1021/acs.cgd.9b00058) http://hdl.handle.net/123456789/2122 (http://hdl.handle.net/123456789/2122) |
| Appears in Collections: | Research Articles (/jspui/handle/123456789/9) |

Files in This Item:

| File | Description | Size | Format | |
|--|-------------|------------|----------------------|---|
| Need to add pdf.odt (/jspui/bitstream/123456789/2122/1/Need%20to%20add%20pdf.odt) | | 8.63 kB | OpenDocument Text | View/Open (/jspui/bitstream/123456789/2122/1/Need%20to%20add%20pdf.odt) |

[Show full item record \(/jspui/handle/123456789/2122?mode=full\)](#)

 [\(/jspui/handle/123456789/2122/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.