



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/5079>


Title:	Design, Synthesis and Characterization of Achiral Unsymmetrical Four-ring based Hockey-stick Shaped Liquid Crystals: Structure-Property relationship
Authors:	Jain, Varsha (/jspui/browse?type=author&value=Jain%2C+Varsha) Kaur, Supreet (/jspui/browse?type=author&value=Kaur%2C+Supreet) Mohiuddin, Golam (/jspui/browse?type=author&value=Mohiuddin%2C+Golam) Pal, Santanu Kumar (/jspui/browse?type=author&value=Pal%2C+Santanu+Kumar)
Keywords:	Design Synthesis Characterization Unsymmetrical Four-ring Hockey-stick
Issue Date:	2021
Publisher:	Taylor & Francis online
Citation:	Liquid Crystals, 49(2), 162–171.
Abstract:	Hockey-stick-shaped liquid crystals are interesting due to their unusual physical properties. Nematic phase observed in these materials have shown promising ferroelectric-like switching, ease of alignment in a planar and homeotropic cell, large flexoelectricity, etc. Polar moieties viz. F, Cl, NO ₂ , etc., in their molecular architecture lead to the development of permanent dipole moment which in turn affects the phase structure and physical properties. Herein, we have designed and synthesised a new series of four-ring-based hockey-stick-shaped molecules having polar Cl group at one end and variable aliphatic chains at another end of the molecular long axis. The four phenyl rings are attached via one ester and two imine linkages. All the compounds exhibited enantiotropic mesomorphism. The lower homologs (1–5 to 1–8) exclusively showed nematic phase whereas other higher homologs either exhibited nematic and/or smectic A phase. Compound 1–5 showed a stable nematic phase with the highest phase range of about 110 °C upon cooling. Detailed X-ray diffraction revealed the internal arrangement of molecules in layers. Density functional theory study confirmed the resultant dipole moment of the molecules (approximately 6 Debye) which in turn cancelled out in the bulk phase due to preferable antiparallel arrangement as a dimer.
Description:	Only IISERM authors are available in the record.
URI:	https://doi.org/10.1080/02678292.2021.1949054 (https://doi.org/10.1080/02678292.2021.1949054) http://hdl.handle.net/123456789/5079 (http://hdl.handle.net/123456789/5079)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format

Need To Add...Full Text_PDF (1) (/jspui/bitstream/123456789/5079/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF%20%281%29)	Only IISERM authors are available in the record.	15.36 kB	Unknown	View
--	--	-------------	---------	----------------------

[Show full item record \(/jspui/handle/123456789/5079?mode=full\)](#)

 [\(/jspui/handle/123456789/5079/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.