



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**  
**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**  
**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4887>

Title:	Geometrical meaning of statistical isotropy of smooth random fields in two dimensions.
Authors:	Yogendran, K. P. (/jspui/browse?type=author&value=Yogendran%2C+K.+P.)
Keywords:	Large scale structure of the Universe Cosmic microwave background
Issue Date:	2021
Publisher:	American Physical Society
Citation:	Physical Review D, 104(12).
Abstract:	<p>We revisit the geometrical meaning of statistical isotropy that is manifest in excursion sets of smooth random fields in two dimensions. Using the contour Minkowski tensor, <math>W_1</math>, as our basic tool we first examine geometrical properties of single structures. For simple closed curves in two dimensions we show that <math>W_1</math> is proportional to the identity matrix if the curve has <math>m</math>-fold symmetry, with <math>m \geq 3</math>. Then we elaborate on how <math>W_1</math> maps any arbitrary shaped simple closed curve to an ellipse that is unique up to translations of its centroid. We also carry out a comparison of the shape parameters, <math>\alpha</math> and <math>\beta</math>, defined using <math>W_1</math>, with the filamentarity parameter defined using two scalar Minkowski functionals—area and contour length. We show that they contain complementary shape information, with <math>W_1</math> containing additional information of orientation of structures. Next, we apply our method to boundaries of excursion sets of random fields and examine what statistical isotropy means for the geometry of the excursion sets. Focusing on Gaussian isotropic fields, and using a seminumerical approach we quantify the effect of finite sampling of the field on the geometry of the excursion sets. In doing so we obtain an analytic expression for <math>\alpha</math> which takes into account the effect of finite sampling. Finally we derive an analytic expression for the ensemble expectation of <math>W_1</math> for Gaussian anisotropic random fields. Our results provide insights that are useful for designing tests of statistical isotropy using cosmological data.</p>
Description:	Only IISER Mohali authors are available in the record.
URI:	<a href="https://doi.org/10.1103/PhysRevD.104.123516">https://doi.org/10.1103/PhysRevD.104.123516</a> ( <a href="https://doi.org/10.1103/PhysRevD.104.123516">https://doi.org/10.1103/PhysRevD.104.123516</a> ) <a href="http://hdl.handle.net/123456789/4887">http://hdl.handle.net/123456789/4887</a> ( <a href="http://hdl.handle.net/123456789/4887">http://hdl.handle.net/123456789/4887</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format	
Need to add pdf (1).odt (/jspui/bitstream/123456789/4887/1/Need%20to%20add%20pdf%20%281%29.odt)		8.63 kB	OpenDocument Text	<a href="#">View/Open (/jspui/bitstream/123456789/4887/1/Need%20to%20add%20pdf%20%281%29.odt)</a>

[Show full item record \(/jspui/handle/123456789/4887?mode=full\)](#)

[Statistics \(/jspui/handle/123456789/4887/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.