



## Library Indian Institute of Science Education and Research Mohali



## DSpace@IISERMohali / Thesis & Dissertation / Master of Science / MS-16

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/3785

Title: Understanding the role of post-synaptic density proteins in the trafficking of group I mGluRs

Authors: Pal, Subhajit.

Post-synaptic Density proteins

ImGluRs

Issue Date:

28-Jul-2021

Publisher:

Keywords:

IISERM

Abstract:

Glutamate is the major excitatory neurotransmitter in the central nervous system (CNS). There are two types of glutamate receptors; ionotropic glutamate receptors (iGluRs) and metabotropic glutamate receptors (mGluRs). mGluRs are divided into three subclasses depending on their sequence similarity, agonist preference and coupled G-proteins. Group I mGluRs are crucial for neuroprotection, neurodevelopment and various kind of synaptic plasticity including learning and memory formation. Aberrations in the glutamate signalling pathway are basis for various neurological and neurodevelopmental disorders like Fragile X syndrome, ischemia, ALS, multiple sclerosis, diabetes, Huntington's disease and Parkinson's disease (Bear MF, Huber KM, Warren ST, 2004). Exact spatio-temporal localization of these receptors at the cell surface is necessary for the normal signalling and functionality of the receptors. Intracellular trafficking of these receptors play critical role in regulating the spatio-temporal localization of these receptors. Although, the agonist-mediated trafficking of group I mGluRs have been studied in some detail, role of the post-synaptic density proteins in the trafficking of these receptors, if any, have not been investigated. Group I mGluRs also undergo internalization through agonist-independent mechanisms. Other than agonists, antagonists are another type of ligand of mGluRs. These molecules bind to the receptor and block their normal activity. There have been few reports where people have shown that antagonist also induces internalization of few GPCRs (B. F. Roettger et al, 1997). But no studies have been done till date to check the role of antagonists in the trafficking of group I mGluRs. The antagonists of these receptors have various therapeutic benefits for various pain states, neuropsychiatric disorders like anxiety and depression, drug addiction and drug withdrawal. The aim of this study was to understand the role of antagonists in the trafficking of group I mGluRs and also to investigate the role

URI: http://hdl.handle.net/123456789/3785

Appears in

MS-16

Collections:

Files in This Item:

File	Description	Size	Format	
MS Thesis MS16030.pdf		2.72 MB	Adobe PDF	View/Open

Show full item record



Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

