



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/5187>

Title:	Measurements of branching fractions and asymmetry parameters of $\Xi^0 c \rightarrow \Lambda K^- \pi^0$, $\Xi^0 c \rightarrow \Sigma^0 K^- \pi^0$, and $\Xi^0 c \rightarrow \Sigma^+ K^- \pi^-$ decays at Belle
Authors:	Patra, Sourav (/jspui/browse?type=author&value=Patra%2C+Sourav)
Keywords:	Branching fraction Charm physics e ⁺ e ⁻ Experiments
Issue Date:	2021
Publisher:	Springer Nature
Citation:	Journal Of High Energy Physics, (6).
Abstract:	Using a data sample of 980 fb ⁻¹ collected with the Belle detector at the KEKB asymmetric-energy e ⁺ e ⁻ collider, we study the processes of $\Xi^0 c \rightarrow \Lambda K^- \pi^0$, $\Xi^0 c \rightarrow \Sigma^0 K^- \pi^0$, and $\Xi^0 c \rightarrow \Sigma^+ K^- \pi^-$ for the first time. The relative branching ratios to the normalization mode of $\Xi^0 c \rightarrow \Xi^- \pi^+$ are measured to be $B(\Xi^0 c \rightarrow \Lambda K^- \pi^0)/B(\Xi^0 c \rightarrow \Xi^- \pi^+) = 0.18 \pm 0.02(\text{stat.}) \pm 0.01(\text{syst.})$, $B(\Xi^0 c \rightarrow \Sigma^0 K^- \pi^0)/B(\Xi^0 c \rightarrow \Xi^- \pi^+) = 0.69 \pm 0.03(\text{stat.}) \pm 0.03(\text{syst.})$, $B(\Xi^0 c \rightarrow \Sigma^+ K^- \pi^-)/B(\Xi^0 c \rightarrow \Xi^- \pi^+) = 0.34 \pm 0.06(\text{stat.}) \pm 0.02(\text{syst.})$, where the uncertainties are statistical and systematic, respectively. We obtain $B(\Xi^0 c \rightarrow \Lambda K^- \pi^0) = (3.3 \pm 0.3(\text{stat.}) \pm 0.2(\text{syst.}) \pm 1.0(\text{ref.})) \times 10^{-3}$, $B(\Xi^0 c \rightarrow \Sigma^0 K^- \pi^0) = (12.4 \pm 0.5(\text{stat.}) \pm 0.5(\text{syst.}) \pm 3.6(\text{ref.})) \times 10^{-3}$, $B(\Xi^0 c \rightarrow \Sigma^+ K^- \pi^-) = (6.1 \pm 1.0(\text{stat.}) \pm 0.4(\text{syst.}) \pm 1.8(\text{ref.})) \times 10^{-3}$, where the uncertainties are statistical, systematic, and from $B(\Xi^0 c \rightarrow \Xi^- \pi^+)$, respectively. The asymmetry parameters $a(\Xi^0 c \rightarrow \Lambda K^- \pi^0)$ and $a(\Xi^0 c \rightarrow \Sigma^+ K^- \pi^-)$ are $0.15 \pm 0.22(\text{stat.}) \pm 0.04(\text{syst.})$ and $-0.52 \pm 0.30(\text{stat.}) \pm 0.02(\text{syst.})$, respectively, where the uncertainties are statistical followed by systematic.
Description:	Only IISER Mohali authors are available in the record.
URI:	http://dx.doi.org/10.1007/JHEP06(2021)160 (http://dx.doi.org/10.1007/JHEP06(2021)160) http://hdl.handle.net/123456789/5187 (http://hdl.handle.net/123456789/5187)
Appears in	Research Articles (/jspui/handle/123456789/9)
Collections:	

Files in This Item:

File	Description	Size	Format	
Need To Add...Full Text_PDF (1) (/jspui/bitstream/123456789/5187/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF%20%281%29)		15.36 kB	Unknown	View

Show full item record (/jspui/handle/123456789/5187?mode=full)

(/jspui/handle/123456789/5187/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

