



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**

**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2759>

Title:	A new bulky iminophosphonamide as an N,N'-chelating ligand: synthesis and structural characterization of heteroleptic group 13 element complexes
Authors:	Prashanth, B. (/jspui/browse?type=author&value=Prashanth%2C+B.) Singh, Sanjay (/jspui/browse?type=author&value=Singh%2C+Sanjay)
Keywords:	Iminophosphonamide Heteroleptic Multinuclear NMR
Issue Date:	2014
Publisher:	Royal Society of Chemistry
Citation:	Dalton Transactions, 43(44), pp.16880-16888.
Abstract:	A sterically demanding iminophosphonamine ligand [(2,6-iPr <sub>2</sub> C <sub>6</sub> H <sub>3</sub> N)P(Ph <sub>2</sub> )(NtBu)]H (LH) and its lithium derivative [(2,6-iPr <sub>2</sub> C <sub>6</sub> H <sub>3</sub> N)P(Ph <sub>2</sub> )(NtBu)](Li·2THF) (1) were used to prepare complexes of group 13 elements. The reaction of LH with AlH <sub>3</sub> ·NMe <sub>2</sub> Et and AlMe <sub>3</sub> respectively, affords [LAH <sub>2</sub> ] <sub>2</sub> (2) and LAIme <sub>2</sub> (3). The lithium derivative 1 when treated with the MCl <sub>3</sub> compound of group 13 yields [(2,6-iPr <sub>2</sub> C <sub>6</sub> H <sub>3</sub> N)P(Ph <sub>2</sub> )(NtBu)]MCl <sub>2</sub> (M = B (4); Al (5); and Ga (6). Compound 3 on reaction with a Lewis acid B(C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> generates the cationic complex [(2,6-iPr <sub>2</sub> C <sub>6</sub> H <sub>3</sub> N)P(Ph <sub>2</sub> )(NtBu)]AlMe <sub>2</sub> <sup>+</sup> [MeB(C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> ] <sup>-</sup> (7) that slowly undergoes rearrangement to yield [(2,6-iPr <sub>2</sub> C <sub>6</sub> H <sub>3</sub> N)P(Ph <sub>2</sub> )(NtBu)]AlMe(C <sub>6</sub> F <sub>5</sub> ) (8) and MeB(C <sub>6</sub> F <sub>5</sub> ) <sub>2</sub> . Compounds 1–8 were characterized using multinuclear NMR, EI-MS and IR techniques and the solid state structure of 1–6 and 8 was elucidated by single crystal X-ray diffraction analyses.
URI:	<a href="https://pubs.rsc.org/en/content/articlelanding/2014/dt/c4dt02637c#!divAbstract">https://pubs.rsc.org/en/content/articlelanding/2014/dt/c4dt02637c#!divAbstract</a> ( <a href="https://pubs.rsc.org/en/content/articlelanding/2014/dt/c4dt02637c#!divAbstract">https://pubs.rsc.org/en/content/articlelanding/2014/dt/c4dt02637c#!divAbstract</a> ) <a href="http://hdl.handle.net/123456789/2759">http://hdl.handle.net/123456789/2759</a> ( <a href="http://hdl.handle.net/123456789/2759">http://hdl.handle.net/123456789/2759</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format	
need to add pdf....odt (/jspui/bitstream/123456789/2759/1/need%20to%20add%20pdf....odt)		8.12 kB	OpenDocument Text	<a href="#">View/Open (/jspui/bitstream/123456789/2759/1/need%20to%20add%20pdf....odt)</a>

Show full item record (/jspui/handle/123456789/2759?mode=full)

(/jspui/handle/123456789/2759/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

