

Library Indian Institute of Science Education and Research Mohali



Viev

DSpace@IISERMohali (/jspui/)

- / Publications of IISER Mohali (/jspui/handle/123456789/4)
- / Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/5185	
Title:	Test of lepton flavor universality and search for lepton flavor violation in B $\to K\ell\ell$ decays
Authors:	Bhardwaj, Vishal (/jspui/browse?type=author&value=Bhardwaj%2C+Vishal) Patra, Sourav (/jspui/browse?type=author&value=Patra%2C+Sourav)
Keywords:	FCNC Interaction Flavour Changing Neutral Currents Flavor physics
Issue Date:	2021
Publisher:	Springer Nature
Citation:	Journal of High Energy Physics, (3).
Abstract:	We present measurements of the branching fractions for the decays B \rightarrow Kµ+µ- and B \rightarrow Ke+e- and their ratio (RK), using a data sample of 711 fb-1 that contains 772 × 106 BB events. The data were collected at the Y(4S) resonance with the Belle detector at the KEKB asymmetric-energy e+e- collider. The ratio RK is measured in five bins of dilepton invariant-mass-squared (q2): q2 ∈ (0.1, 4.0), (4.00, 8.12), (1.0, 6.0), (10.2, 12.8) and (> 14.18) GeV2/c4, along with the whole q2 region. The RK value for q2 ∈ (1.0, 6.0) GeV2/c4 is 1.03+0.28-0.24 ± 0.01. The first ansecond uncertainties listed are statistical and systematic, respectively. All results for RK are consistent with Standard Model predictions. We also measure CP-averaged isospin asymmetries in the same q2 bins. The results are consistent with a null asymmetry, with the largest difference of 2.6 standard deviations occurring for the q2 ∈ (1.0, 6.0) GeV2/c4 bin in the mode with muon final states. The measured differential branching fractions, dB/dq2, are consistent with theoretical predictions for charged B decays, while the corresponding values are below the expectations for neutral B decays. We have also searched for lepton-flavor-violating B \rightarrow Kµ±e \mp decays and set 90% confidence-level upper limits on the branching fraction in the range of 10-8 for B+ \rightarrow K+µ±e \mp , and B0 \rightarrow K0µ±e \mp modes.
Description:	Only IISER Mohali authors are available in the record.

Description.	Offig Hotel Worldh authors are available in the record.

URI: https://dx.doi.org/10.1007/jhep03(2021)105 (https://dx.doi.org/10.1007/jhep03(2021)105)

http://hdl.handle.net/123456789/5185 (http://hdl.handle.net/123456789/5185)

Appears in Research Articles (/jspui/handle/123456789/9) Collections:

Files in This Item:

 File
 Description
 Size
 Format

 Need To Add...Full Text_PDF (1)
 15.36
 Unknown (/jspui/bitstream/123456789/5185/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF%20%281%29)
 kB

Show full item record (/jspui/handle/123456789/5185?mode=full)

. (/jspui/handle/123456789/5185/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.