



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2856>


Title:	Diastereoselective Construction of 3-Aminooxindoles with Adjacent Stereocenters: Stereocontrolled Addition of γ -Substituted Allylindiums to Isatin Ketimines
Authors:	Aslam, N.A. (/jspui/browse?type=author&value=Aslam%2C+N.A.) Babu, S.A. (/jspui/browse?type=author&value=Babu%2C+S.A.) Rani, Soniya (/jspui/browse?type=author&value=Rani%2C+Soniya) Mahajan, Shivam (/jspui/browse?type=author&value=Mahajan%2C+Shivam) Solanki, Jagmohan (/jspui/browse?type=author&value=Solanki%2C+Jagmohan)
Keywords:	diastereoselective construction 3-Aminooxindoles γ -Substituted Allylindiums
Issue Date:	2015
Publisher:	WILEY-VCH Verlag GmbH
Citation:	European Journal of Organic Chemistry, 205(19)
Abstract:	The diastereoselective construction of 3-allyl-3-aminooxindoles that have two adjacent stereocenters has been achieved by the In-promoted Barbier-type addition of γ -substituted allylic halides to the C=N bond of isatin ketimines. The reactions of cinnamyl-, crotyl-, and geranylium compounds with isatin ketimines proceeded in either aqueous or alcohol solution. The addition of a cyclohexenylindium species to an isatin ketimine was carried out in N,N-dimethylformamide (DMF), and the addition of ethyl 4-bromocrotonate to an isatin ketimine in EtOH gave oxindole-based β -amino acid scaffolds. In all of these processes, the reaction conditions were screened to obtain the respective 3-allyl-3-aminooxindoles with very high stereoselectivity. In addition, plausible TS models are proposed, and representative synthetic transformations were carried out by using the oxindole-based β -amino acid scaffolds. Furthermore, the stereochemistry of representative compounds were unequivocally established by single-crystal X-ray structure analysis. A synthetic protocol for the diastereoselective indium-mediated Barbier-type C-C bond formation has been developed. The addition of γ -substituted allylic reagents to the C=N bond of isatin ketimines was employed for the synthesis of several new 3-aminooxindoles that have two contiguous stereocenters.
Description:	Only IISERM authors are available in the record.
URI:	https://chemistry-europe.onlinelibrary.wiley.com/doi/full/10.1002/ejoc.201500340 (https://chemistry-europe.onlinelibrary.wiley.com/doi/full/10.1002/ejoc.201500340) http://hdl.handle.net/123456789/2856 (http://hdl.handle.net/123456789/2856)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.ott (/jspui/bitstream/123456789/2856/1/Need%20to%20add%20pdf.ott)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/2856/1/Need%20to%20add%20pdf.ott\)](#)

Show full item record (</jspui/handle/123456789/2856?mode=full>)

 (</jspui/handle/123456789/2856/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.