

Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

- / Publications of IISER Mohali (/jspui/handle/123456789/4)
- / Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/2422					
Title:	Classification of flat connected quandles				
Authors:	Singh, Mahender (/jspui/browse?type=author&value=Singh%2C+Mahender)				
Keywords:	Automorphism of quandle				
	Central automorphism				
	Connected quandle				
	Flat quandle				
Issue Date:	2016				
Publisher:	World Scientific				
Citation:	Journal of Knot Theory and its Ramifications,25(13).				
Abstract:	Let A be an additive abelian group. Then the binary operation $a*b=2b-a$ gives a quandle structure on A, denoted by T(A), and called the Takasaki quandle of A. Viewing quandles as generalization of Riemannian symmetric spaces, Ishihara and Tamaru [Flat connected finite quandles, to appear in Proc. Amer. Math. Soc. (2016)] introduced flat quandles, and classified quandles which are finite, flat and connected. In this note, we classify all flat connected quandles. More precisely, we prove that a quandle X is flat and connected if and only if $X \cong T(A)$, where A is a 2-divisible group.				
URI:	https://www.worldscientific.com/doi/10.1142/S0218216516500711				
	(https://www.worldscientific.com/doi/10.1142/S0218216516500711)				
	http://hdl.handle.net/123456789/2422 (http://hdl.handle.net/123456789/2422)				
Appears in	Research Articles (/jspui/handle/123456789/9)				

Files in This Item:

Collections:

File	Description	Size	Format	
Need to add pdf.odt (/jspui/bitstream/123456789/2422/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text	View/Open (/jspui/bitstream/12345

Show full item record (/jspui/handle/123456789/2422?mode=full)

■ (/jspui/handle/123456789/2422/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.