



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali / Thesis & Dissertation / Doctor of Philosophy (PhD) / PhD-2013

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4033>

Title:	Skymions and antiskymions in spin-orbit modified double exchange models
Authors:	Kathyat, Deepak Singh
Keywords:	Skymions antiskymions
Issue Date:	Jul-2021
Publisher:	IISER Mohali
Abstract:	<p>It has been understood that the conventional electronics based devices will not be able to meet the ever increasing requirements for data storage and processing of the modern world. Among others, the idea of utilizing the spin degree of freedom associated with electrons has been considered as a potential alternative. This has given rise to a new research field, popularly known as spintronics, wherein electron's spin is used as the carrier of information for device functionalities. While it is not easy to detect and utilize spin of a single electron in materials containing a large number of them, certain magnetic configurations show a desired stability and possible control via external electric or magnetic field. Therefore, search for magnetic materials supporting certain stable magnetic textures has become a key theme of research in recent years. Topologically protected magnetic textures, such as skymions and anti-skymions are of special importance due to their stability. Such textures have been discovered in chiral magnets and in thin films of a variety of magnetic metals. The fundamental physics associated with formation of magnetic skymions has fascinated researchers since the discovery of these topological textures. The current approach to understand these intriguing textures is via spin models consisting of Dzyaloshinskii-Moriya (DM) interactions or frustrating long range interactions. In this thesis, we present a microscopic mechanism for skymion and antiskymion formation in metals that emerges from electronic itinerancy. We derive and study a microscopic spin Hamiltonian on a lattice for double exchange metals modified by the Rashba and Dresselhaus spin orbit coupling (SOC). In our model, anisotropic interactions of the Dzyaloshinskii-Moriya (DM) and pseudo dipolar form emerge naturally in addition to the standard isotropic term. We present phase diagram of the effective spin Hamiltonian which has very interesting ground states like classical spin liquid state using large scale Monte Carlo simulations. We show that in presence of Zeeman field the mechanism we propose not only provides an accurate microscopic understanding of existence of skymions, but also explains key features in small angle neutron scattering (SANS) and Lorentz transmission electron microscopy (LTEM) data on thin films of MnSi-type B20 metals and transition metals and their alloys. We identify hexagonal and square lattice arrangements of skymions in two different regimes of the parameter space. Sparse skymions emerge at finite temperatures as excitations of the ferromagnetic phase. Further, the skymion states are characterized as topological metals via explicit calculations of Bott index and Hall conductivity. Local density of states (LDOS) display characteristic oscillations arising from a combination of confinement effect and gauge-field induced Landau level physics. These unique features serve as testable predictions for the presence of the new mechanism of skymion formation in real materials. The discovery of a new mechanism based on two celebrated physics concepts not only fills a major conceptual void in the current understanding of skymions and antiskymions in metals, but also opens a new route for tuning the size, density and stability of skymions in magnetic metals. We also emphasize the importance of a consistent treatment of spin-orbit coupling for calculating electronic properties of metals hosting unconventional magnetic textures such as skymions. Finally, we provide a clear understanding of how Neel-type skymions, Bloch-type skymions and the corresponding antiskymions are related with one another within a simple lattice model. We also emphasize the role played by electron itinerancy in deciding the type of skymion textures in a metal. These features are completely missed in a spin-only model written without reference to a starting microscopic model.</p>
URI:	http://hdl.handle.net/123456789/4033
Appears in Collections:	PhD-2013

Files in This Item:

File	Description	Size	Format	
thesis Deepak Singh Kathyat PH13003.pdf		32.48 MB	Adobe PDF	View/Open

Show full item record



Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

Admin Tools

[Edit...](#)

[Export Item](#)

Export (migrate) Item

Export metadata



Customized & Implemented by - [Jivesna Tech](#)