



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Thesis & Dissertation (/jspui/handle/123456789/1)

/ Doctor of Philosophy (PhD) (/jspui/handle/123456789/268)

/ PhD-2012 (/jspui/handle/123456789/955)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1217>


Title:	Understanding the gene regulatory network underlying zebrafish retina regeneration
Authors:	Kaur, Simran (/jspui/browse?type=author&value=Kaur%2C+Simran)
Keywords:	Biology RNA isolation Microscopy Fluorescence assisted cell sorting Retina regeneration
Issue Date:	1-Oct-2019
Publisher:	IISERM
Abstract:	<p>Visual impairment is one of the major problems faced by humanity. Despite the fact that the notion of retina regeneration seems implausible, unlike humans, there are vertebrates such as piscine and amphibians that have remarkable retina regeneration capabilities. Such organisms serve as a great tool to study retina regeneration, with zebrafish being a prominent example. Zebrafish, possess a significant regenerative response and with a retinal architecture similar to mammals, has emerged as the baseline model for studying retina regeneration. Upon retinal injury in zebrafish, Muller glia cells reprograms themselves to a multipotent stem cell-like state (Muller glial derived progenitor cells - MGPCs), which then proliferate further and give rise to various retinal cell types. Although many molecular signaling pathways and transcription factors are known to play an essential role during retina regeneration, one of the most crucial developmental signaling pathways, i.e., Sonic hedgehog (Shh) signaling pathway remained under-explored. In this study, we show that the induction of Shh signaling is inevitable for zebrafish retina regeneration. Further, we report on a let-7 microRNA-mediated stringent translational regulation of the components of Shh signaling such as sonic hedgehog a (shha), sonic hedgehog b (shhb), patched1 (ptch1) and smoothened (smo). We also show the direct necessity of Shh signaling for the induction of pivotal transcription factor genes like ascl1a, protein, lin28a. Interestingly, Matrix zic2b foxn4 and metalloproteinase-9 a RNA (Mmp9) binding causes the initiation of Shh signaling for reprogramming the Muller glial cells to MGPCs. Further investigation showed the repressive role of hairy enhancer of split-related (Her4.1, an effector of Notch signaling) in controlling the expression of mmp9, precisely in proliferating Muller glial cells. We believe that these novel findings pave the way to design new therapeutic strategies to treat retinal diseases in humans.</p>
URI:	IISERM (IISERM) http://hdl.handle.net/123456789/1217 (http://hdl.handle.net/123456789/1217)

Appears in PhD-2012 (/jspui/handle/123456789/955)
Collections:

Files in This Item:

File	Description	Size	Format	
PH12146.pdf (/jspui/bitstream/123456789/1217/4/PH12146.pdf)		72.42 MB	Adobe PDF	View/Open (/jspui/bitstream/123456789/1217/4/PH12146.pdf)

[Show full item record \(/jspui/handle/123456789/1217?mode=full\)](/jspui/handle/123456789/1217?mode=full)

 [\(/jspui/handle/123456789/1217/statistics\)](/jspui/handle/123456789/1217/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

Admin Tools

[Edit...](#)

[Export Item](#)

[Export \(migrate\) Item](#)

[Export metadata](#)