



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**

**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4583>


Title:	Spectral States of OJ 287 blazar from Multiwavelength Observations with AstroSat
Authors:	Singh, K P (/jspui/browse?type=author&value=Singh%2C+K+P)
Keywords:	Spectral States Multiwavelength AstroSat
Issue Date:	2022
Publisher:	Oxford Academic
Citation:	Monthly Notices of the Royal Astronomical Society, 509(2), 2696-2706.
Abstract:	We present AstroSat soft X-ray, near-UV (NUV), and far-UV (FUV) observations of a blazar, OJ 287, carried out in 2017, 2018, and 2020. The simultaneous observations with NuSTAR in 2017 provide a broad-band look encompassing NUV, FUV, soft, and hard X-rays. Captured in three different broad-band spectral states in three observations, the X-ray spectrum is found to be the hardest during 2018, while the high-energy-end of the simultaneous optical-FUV spectrum shows a steepening that is modelled with a broken power-law spectrum. The spectral energy distribution (SED) in 2017 shows a relatively flatter optical-FUV and soft X-ray spectra, implying an additional emission component. The 2020 optical-FUV spectrum is harder than in 2017 and 2018, with an extremely soft X-ray spectrum and a hardening above $\sim 1$ GeV, similar to the SEDs of High-energy-peaked BL Lac objects (HBL), thereby establishing that this additional emission component has HBL-like properties. The AstroSat multiwavelength observations trace the spectral evolution from the end-phase of the HBL component in 2017 to its disappearance in 2018 followed by its revival in 2020. A single zone leptonic model reproduces the 2018 broad-band spectrum, while the 2017 and 2020 SEDs require an additional HBL-like emitting zone. The spectral evolution of the high-energy-end of optical-UV spectrum, revealed by the FUV observations in 2017 and 2018, strongly suggests that X-ray spectral changes in the normal broad-band spectral state of OJ 287 are primarily due to the evolution of the optical-UV synchrotron spectrum.
Description:	Only IISER Mohali authors are available in the record.
URI:	<a href="https://doi.org/10.1093/mnras/stab3161">https://doi.org/10.1093/mnras/stab3161</a> ( <a href="https://doi.org/10.1093/mnras/stab3161">https://doi.org/10.1093/mnras/stab3161</a> ) <a href="http://hdl.handle.net/123456789/4583">http://hdl.handle.net/123456789/4583</a> ( <a href="http://hdl.handle.net/123456789/4583">http://hdl.handle.net/123456789/4583</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need To Add...Full Text_PDF. (/jspui/bitstream/123456789/4583/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF.)		15.36 kB	Unknown

[View/Open \(/jspui/\)](#)

Show full item record (</jspui/handle/123456789/4583?mode=full>)

 (</jspui/handle/123456789/4583/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.