



| | |
|---|---|
| Title: | (2 S)-2-[(Phenylsulfinyl)methyl]pyrrolidine-catalyzed efficient stereoselective michael addition of cyclohexanone and cyclopentanone to nitroolefins |
| Authors: | Khullar, S. (/jspui/browse?type=author&value=Khullar%2C+S.)<br>Mandal, S.K. (/jspui/browse?type=author&value=Mandal%2C+S.K.) |
| Keywords: | Michael addition reaction<br>Organocatalyst<br>Amino sulfoxide<br>Nitroolefins |
| Issue Date: | 2013 |
| Publisher: | Thieme |
| Citation: | Synthesis (Germany), 45(10), pp.1406-1413. |
| Abstract: | (2S)-2-[(Phenylsulfinyl)methyl]pyrrolidine, derived from l-proline, has been demonstrated as an efficient organocatalyst for the asymmetric Michael addition of cyclohexanone and cyclopentanone to β-nitrostyrenes. This pyrrolidine-based catalyst bearing a sulfoxide moiety was used to synthesize various γ-nitro carbonyl compounds in high yield (up to 97%) with excellent stereoselectivity (up to >99:1 dr and >99% ee) without the use of any additive. |
| Description: | Only IISERM authors are available in the record. |
| URI: | https://www.thieme-connect.com/products/ejournals/abstract/10.1055/s-0032-1316917 (https://www.thieme-connect.com/products/ejournals/abstract/10.1055/s-0032-1316917)<br>http://hdl.handle.net/123456789/2931 (http://hdl.handle.net/123456789/2931) |
| Appears in Collections: | Research Articles (/jspui/handle/123456789/9) |

Files in This Item:

| File | Description | Size | Format | |
|---|---|---|---|---|
| Need to add pdf.odt (/jspui/bitstream/123456789/2931/1/Need%20to%20add%20pdf.odt) | | 8.63 kB | OpenDocument Text | View/Open (/jspui/bitstream/12345 |

Show full item record (/jspui/handle/123456789/2931?mode=full)
(/jspui/handle/123456789/2931/statistics)