



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2841>

Title:	Appearance of annular ring-like intermediates during amyloid fibril formation from human serum albumin
Authors:	Arya, S. (/jspui/browse?type=author&value=Arya%2C+S.) Kumari, Arpana (/jspui/browse?type=author&value=Kumari%2C+Arpana) Dalal, Vijit (/jspui/browse?type=author&value=Dalal%2C+Vijit) Bhattacharya, M. (/jspui/browse?type=author&value=Bhattacharya%2C+M.) Mukhopadhyay, S. (/jspui/browse?type=author&value=Mukhopadhyay%2C+S.)
Keywords:	Annular ring-like Amyloid fibril Human serum
Issue Date:	2015
Publisher:	Royal Society of Chemistry
Citation:	Physical Chemistry Chemical Physics, 17(35)
Abstract:	The self-assembly of proteins triggered by a conformational switch into highly ordered β -sheet rich amyloid fibrils has captivated burgeoning interest in recent years due to the involvement of amyloids in a variety of human diseases and a diverse range of biological functions. Here, we have investigated the mechanism of fibrillogenesis of human serum albumin (HSA), an all- α -helical protein, using an array of biophysical tools that include steady-state as well as time-resolved fluorescence, circular dichroism and Raman spectroscopy in conjunction with atomic force microscopy (AFM). Investigations into the temporal evolution of nanoscale morphology using AFM revealed the presence of ring-like intermediates that subsequently transformed into worm-like fibrils presumably by a ring-opening mechanism. Additionally, a multitude of morphologically-diverse oligomers were observed on the pathway to amyloid formation. Kinetic analysis using multiple structural probes in-tandem indicated that HSA amyloid assembly is a concerted process encompassing a major structural change that is primarily mediated by hydrophobic interactions between thermally-induced disordered segments originating in various domains. A slower growth kinetics of aggregates suggested that the protein structural reorganization is a prerequisite for fibril formation. Moreover, time-dependent Raman spectroscopic studies of HSA aggregation provided key molecular insights into the conformational transitions occurring within the protein amide backbone and at the residue-specific level. Our data revealed the emergence of conformationally-diverse disulfides as a consequence of structural reorganization and sequestration of tyrosines into the hydrophobic amyloid core comprising antiparallel cross β -sheets
URI:	https://pubs.rsc.org/en/content/articlelanding/2015/cp/c5cp03782d#!divAbstract (https://pubs.rsc.org/en/content/articlelanding/2015/cp/c5cp03782d#!divAbstract) http://hdl.handle.net/123456789/2841 (http://hdl.handle.net/123456789/2841)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:


File	Description	Size	Format
------	-------------	------	--------

Need to add pdf.odt
(/jspui/bitstream/123456789/2841/1/Need%20to%20add%20pdf.odt)

8.63 OpenDocument
kB Text

[View/Open \(/jspui/bitstream/123456789/2841/1/Need%20to%20add%20pdf.odt\)](#)

[Show full item record \(/jspui/handle/123456789/2841?mode=full\)](#)

 [\(/jspui/handle/123456789/2841/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.