



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2236>

Title:	Molecular and Functional Insights into the Regulation of d-Galactonate Metabolism by the Transcriptional Regulator DgoR in Escherichia coli
Authors:	Singh, Bhupinder (/jspui/browse?type=author&value=Singh%2C+Bhupinder) Arya, G. (/jspui/browse?type=author&value=Arya%2C+G.) Kundu, Neeladrita (/jspui/browse?type=author&value=Kundu%2C+Neeladrita) Sangwan, A. (/jspui/browse?type=author&value=Sangwan%2C+A.) Nongthombam, Shachikanta (/jspui/browse?type=author&value=Nongthombam%2C+Shachikanta) Chaba, Rachna (/jspui/browse?type=author&value=Chaba%2C+Rachna)
Keywords:	GntR family Carbohydrate metabolism Dgo operon
Issue Date:	2019
Publisher:	American Society for Microbiology
Citation:	Journal of Bacteriology, 201(4).
Abstract:	d-Galactonate, an aldonic sugar acid, is used as a carbon source by Escherichia coli, and the structural dgo genes involved in its metabolism have previously been investigated. Here, using genetic, biochemical and bioinformatics approaches, we present the first detailed molecular and functional insights into the regulation of d-galactonate metabolism in E. coli K-12 by the transcriptional regulator DgoR. We found that dgoR deletion accelerates the growth of E. coli in d-galactonate concomitant with the strong constitutive expression of dgo genes. In the dgo locus, sequence upstream of dgoR alone harbors the d-galactonate-inducible promoter that likely drives the expression of all dgo genes. DgoR exerts repression on the dgo operon by binding two inverted repeats overlapping the dgo promoter. Binding of d-galactonate induces a conformational change in DgoR to derepress the dgo operon. The findings from our work firmly place DgoR in the GntR family of transcriptional regulators: DgoR binds an operator sequence [5'-TTGTA(G/C)TACA(A/T)-3'] matching the signature of GntR family members that recognize inverted repeats [5'-(N)yGT(N)xAC(N)y-3', where x and y indicate the number of nucleotides, which varies], and it shares critical protein-DNA contacts. We also identified features in DgoR that are otherwise less conserved in the GntR family. Recently, missense mutations in dgoR were recovered in a natural E. coli isolate adapted to the mammalian gut. Our results show these mutants to be DNA binding defective, emphasizing that mutations in the dgo-regulatory elements are selected in the host to allow simultaneous induction of dgo genes. The present study sets the basis to explore the regulation of dgo genes in additional enterobacterial strains where they have been implicated in host-bacterium interactions.
URI:	https://jb.asm.org/content/201/4/e00281-18 (https://jb.asm.org/content/201/4/e00281-18) http://hdl.handle.net/123456789/2236 (http://hdl.handle.net/123456789/2236)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format	
Need to add pdf.odt (/jspui/bitstream/123456789/2236/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text	View/Open (/jspui/bitstream/123456789/2236/1/Need%20to%20add%20pdf.odt)

[Show full item record \(/jspui/handle/123456789/2236?mode=full\)](#)

 [\(/jspui/handle/123456789/2236/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.