



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Thesis & Dissertation (/jspui/handle/123456789/1)
/ Doctor of Philosophy (PhD) (/jspui/handle/123456789/268)
/ PhD-2011 (/jspui/handle/123456789/954)


Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1190>

Title:	Understanding synaptic function and RNA splicing using <i>C. elegans</i> as a model system
Authors:	Sharma, Pallavi (/jspui/browse?type=author&value=Sharma%2C+Pallavi)
Keywords:	Biology Microscopy RNA splicing using <i>C. elegans</i> Neuromuscular Development Quantitative PCR
Issue Date:	28-Sep-2019
Publisher:	IISERM
Abstract:	<p>Caenorhabditis elegans, a small nematode with its genome being sequenced, well- defined nervous system and pre determined cell fate has been serving the scientific community to study various cellular and biological processes. More the 60% of <i>C. elegans</i> genes have human counterparts, which makes it an excellent model system to study diverse biological processes such as neurobiology, aging, apoptosis, gene regulation and developmental biology. In my work I have used <i>C. elegans</i> to understand two different phenomena; synaptic functioning and RNA splicing. In first part of my talk, I will discuss characterization of the function of a claudin- like protein, HPO-30, and its role in maintaining the levamisole sensitive nicotinic acetylcholine receptors (LACHRs) at the neuromuscular junction (NMJ). Using pharmacological and electrophysiological approaches we establish that in hpo-30 mutants, the LACHR levels are compromised at the NMJ. HPO-30 localizes at the NMJ and shows genetic and physical interaction with the LACHRs. Finally, we show that HPO-30 functions through another cell adhesion molecule neuroligin (NLG-1) to maintain the postsynaptic receptor levels. The second part of my talk involves understanding the function of a ubiquitin like protein, Hub1 in <i>C. elegans</i>. In this work, we identify the interaction of CeHub1 with splicosomal protein CeSNU66. Further, using microarray analysis we assay for splicing specific function of Hub1 in <i>C. elegans</i> and establish that Hub1 function is quite conserved across species and plays an important role in mRNA splicing in <i>C.elegans</i>.</p>
URI:	IISERM (IISERM) http://hdl.handle.net/123456789/1190 (http://hdl.handle.net/123456789/1190)
Appears in Collections:	PhD-2011 (/jspui/handle/123456789/954)

Files in This Item:

File	Description	Size	Format	
PH11099.pdf (/jspui/bitstream/123456789/1190/3/PH11099.pdf)		5.4 MB	Adobe PDF	View/Open (/jspui/bitstream/123456789/1190/3/I

[Show full item record \(/jspui/handle/123456789/1190?mode=full\)](#)

 [\(/jspui/handle/123456789/1190/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

Admin Tools

[Edit...](#)

[Export Item](#)

[Export \(migrate\) Item](#)

[Export metadata](#)