



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)


Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2977>

Title:	Understanding of the Weak Intermolecular Interactions Involving Halogens in Substituted N-Benzylideneanilines: Insights from Structural and Computational Perspectives
Authors:	Kaur, Gurpreet (/jspui/browse?type=author&value=Kaur%2C+Gurpreet) Choudhury, A.R. (/jspui/browse?type=author&value=Choudhury%2C+A.R.)
Keywords:	N-Benzylideneanilines Halogens Hydrogen bonds
Issue Date:	2014
Publisher:	American Chemical Society
Citation:	Crystal Growth and Design,14(4), pp.1600-1616.
Abstract:	The C-F group, which is found in a large number of small organic molecules and drugs available in the market, has still not been fully understood in terms of the strength and directionality of the interactions offered by this group in guiding the formation of crystal lattices. In this manuscript, we have tried to understand the role played by the C-F group, using a model system of N-benzylideneanilines, on which we have previously done a systematic study with fluorine as a substituent on both rings. The effect on the packing of these molecules by replacing one of the fluorine atoms by either Cl or Br has been comprehended in this manuscript. It was observed that the features of the difluorinated analogues remained intact when the noninteracting fluorine atom was replaced by Cl or Br, while with the replacement of the interacting fluorine by Cl or Br, completely different packing characteristics were found to be developed. To quantify the strength of the interactions offered by "organic fluorine", stabilization energies of the dimers (which has been found to interact through the C-H...F hydrogen bond) have been calculated by Gaussian 09 at the MP2 level using a 6-31+G* basis set. These values were found to be between -0.3 and -6.0 kcal/mol. To study the topological properties of the interacting molecular pair, AIM calculations have also been done using AIM2000. In the studied dimers, the existence of bond critical points (BCPs) at the C-H...F hydrogen bond have always been seen and Laplacian at those BCPs has also been found to be positive, which is clearly an indication of a closed shell type of interaction between the C-H and F-C groups.
URI:	https://pubs.acs.org/doi/10.1021/cg401573d (https://pubs.acs.org/doi/10.1021/cg401573d) http://hdl.handle.net/123456789/2977 (http://hdl.handle.net/123456789/2977)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format	
need to add pdf....odt (/jspui/bitstream/123456789/2977/1/need%20to%20add%20pdf....odt)		8.12 kB	OpenDocument Text	View/Open (/jspui/bitstream/123456789/2977/1/need%20to%20add%20pdf....odt)

Show full item record (</jspui/handle/123456789/2977?mode=full>)

 (</jspui/handle/123456789/2977/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.