



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali / Thesis & Dissertation / Master of Science / MS-19

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/5796>

Title:	Elucidating the role of CRP32 (Cysteine-rich peptide 32) and LRR-RLK3 (leucine-rich repeat receptor kinase3) under salinity stress in Arabidopsis thaliana
Authors:	Swapna, Vadthya
Keywords:	ecosystem sodium chloride
Issue Date:	May-2024
Publisher:	IISER Mohali
Abstract:	Salt, consisting of essential sodium and chloride ions, serves as a critical micronutrient, and is vital for robust plant growth and development, with implications extending to higher trophic levels. Sodium and chloride ions are pivotal for maintaining osmotic equilibrium within plant cells, and their deficiency can disrupt water movement, leading to dehydration and growth impairment. Conversely, excess salt, particularly sodium chloride (NaCl), induces osmotic stress in plants and disrupts electrolyte balance in animals, emphasizing the necessity of understanding stress adaptation pathways for sustainable agriculture and human health. Plants have evolved sophisticated strategies, including ion exclusion, osmotic adjustment, and antioxidant defense mechanisms, to mitigate salt-induced damage and ensure cellular homeostasis, enabling survival even in saline environments. A particularly intriguing aspect is the role of Cysteine-Rich Peptides (CRPs) in stress-mediated signaling, orchestrating salt nutrition homeostasis and illuminating the intricate molecular choreography of resilience in adverse conditions, thereby offering promising avenues for enhancing crop productivity and ecosystem resilience.
URI:	http://hdl.handle.net/123456789/5796
Appears in Collections:	MS-19

Files in This Item:

File	Description	Size	Format	
embargo period.pdf		6.04 kB	Adobe PDF	View/Open

Show full item record



Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.