



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**

**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/5080>


Title:	Deciphering supramolecular isomerization in coordination polymers: connected molecular squares vs. fused hexagons
Authors:	Gupta, Vijaya (/jspui/browse?type=author&value=Gupta%2C+Vijaya) Laha, Biswajit (/jspui/browse?type=author&value=Laha%2C+Biswajit) Khullar, Sadhika (/jspui/browse?type=author&value=Khullar%2C+Sadhika) Mandal, Sanjay K. (/jspui/browse?type=author&value=Mandal%2C+Sanjay+K.)
Keywords:	Deciphering supramolecular isomerization coordination polymers
Issue Date:	2021
Publisher:	Dalton Transaction
Citation:	Dalton Transactions, 50(6),2221–2232.
Abstract:	The self-assembly of Mn(II), bis(tridentate) ligands and bent dicarboxylate linkers under ambient conditions has been exploited to generate a series of 1D coordination polymers in good yields. For a set of seven compounds, structural isomerization of these architectures is demonstrated through the variation of length and nature of the spacer between the tridentate capping sites of the bis(tridentate) ligands, such as tpbn (N,N',N'',N'''-tetrakis-(2-pyridylmethyl)-1,4-diaminobutane), tpxn (N,N',N'',N'''-tetrakis-(2-pyridylmethyl)-1,6-diaminohexane), and tpxn (N,N',N'',N'''-tetrakis-(2-pyridylmethyl)-xylylamine) or by varying the bent dicarboxylate linker 4,4'-(dimethylsilanediyl)bis-benzoic acid (H2L1) or 4,4'-oxybis-benzoic acid (H2L2). These compounds have been structurally characterized by single-crystal and powder X-ray diffraction, FTIR, and thermogravimetric and elemental analyses. This study reveals that the supramolecular structural variation can be precisely controlled either by a judicious selection of reaction conditions or linker/ligand combinations. For example, the self-assembly of Mn(II), tpbn and H2L1 in DMF/EtOH/water affords a mixture of products (1 and 1a) while changing the solvent combination to EtOH/water results in the generation of a single isomer (1a) in a highly selective manner. On the other hand, for the Mn(II)-tpxn system, different structural isomers have been isolated by varying the dicarboxylates, H2L1 and H2L2 (2vs.5). Similarly, for the Mn(II)-H2L2 system, a variation in the spacer chain length of bis(tridentate) ligands, tpbn and tpxn resulted in the formation of different structural isomers (4vs.5).
Description:	Only IISER authors are available in the record.
URI:	<a href="https://doi.org/10.1039/d0dt04196c">https://doi.org/10.1039/d0dt04196c</a> ( <a href="https://doi.org/10.1039/d0dt04196c">https://doi.org/10.1039/d0dt04196c</a> ) <a href="http://hdl.handle.net/123456789/5080">http://hdl.handle.net/123456789/5080</a> ( <a href="http://hdl.handle.net/123456789/5080">http://hdl.handle.net/123456789/5080</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format

Need To Add...Full Text_PDF (1) (/jspui/bitstream/123456789/5080/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF%20%281%29)	Only IISERM authors are available in the record.	15.36 kB	Unknown	<a href="#">View</a>
--	--	-------------	---------	----------------------

[Show full item record \(/jspui/handle/123456789/5080?mode=full\)](/jspui/handle/123456789/5080?mode=full)

 [\(/jspui/handle/123456789/5080/statistics\)](/jspui/handle/123456789/5080/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.