



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**

**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2294>

Title:	A robust and water-stable two-fold interpenetrated metal–organic framework containing both rigid tetrapodal carboxylate and rigid bifunctional nitrogen linkers exhibiting selective CO <sub>2</sub> capture†
Authors:	Gupta, Vijay (/jspui/browse?type=author&value=Gupta%2C+Vijay) Mandal, S.K. (/jspui/browse?type=author&value=Mandal%2C+S.K.)
Keywords:	Microporous Metal–organic framework Self-assembly
Issue Date:	2019
Publisher:	American Chemical Society
Citation:	Dalton Transactions, 48(2), pp.415-425.
Abstract:	A microporous metal–organic framework, {[Co <sub>2</sub> (4,4'-bpy)(L)]·H <sub>2</sub> O·0.5(DMF)} <sub>n</sub> (1), was obtained from the self-assembly of cobalt(II) nitrate hexahydrate, rigid tetrapodal carboxylic acid 4,4',4'',4'''-silanetetrayltetrabenzoic acid (H <sub>4</sub> L) and a rigid bifunctional linker 4,4'-bipyridine (4,4'-bpy) under solvothermal conditions. It was characterized by Fourier transform infrared spectroscopy, thermogravimetric analysis, elemental analysis, and powder and single crystal X-ray diffraction. Its single crystal structure reveals the presence of a Co(II)-paddle wheel core as the SBU, which is extended to form a doubly interpenetrated 3D framework with a (4,6)-connected sqc422-type uninodal net topology with the Schläfli point symbol {42·510·72·8}{42·54}. There are two types of open channels in this framework (rhombic and trigonal), which run along all three axes. Its thermal and chemical stabilities were established based on thermogravimetric analysis and in situ variable temperature powder X-ray diffraction. The activated framework (lattice solvent free) of 1 exhibits modest uptake of CO <sub>2</sub> (53.8 and 36.4 cm <sup>3</sup> g <sup>-1</sup> at 273 and 298 K at 1 bar pressure, respectively), with moderately high selectivities for CO <sub>2</sub> /N <sub>2</sub> and CO <sub>2</sub> /CH <sub>4</sub> gas separation under ambient conditions (298 and 273 K under 1 bar pressure).
URI:	<a href="https://pubs.rsc.org/en/content/articlelanding/2019/dt/c8dt03844a#!divAbstract">https://pubs.rsc.org/en/content/articlelanding/2019/dt/c8dt03844a#!divAbstract</a> ( <a href="https://pubs.rsc.org/en/content/articlelanding/2019/dt/c8dt03844a#!divAbstract">https://pubs.rsc.org/en/content/articlelanding/2019/dt/c8dt03844a#!divAbstract</a> ) <a href="http://hdl.handle.net/123456789/2294">http://hdl.handle.net/123456789/2294</a> ( <a href="http://hdl.handle.net/123456789/2294">http://hdl.handle.net/123456789/2294</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/2294/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/2294/1/Need%20to%20add%20pdf.odt\)](#)

[Show full item record \(/jspui/handle/123456789/2294?mode=full\)](#)

[Statistics \(/jspui/handle/123456789/2294/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.