



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2705>

Title: Disulphide bond restrains the C-terminal region of thermostable direct hemolysin during folding to promote oligomerization

Authors: Kundu, Nidhi (/jspui/browse?type=author&value=Kundu%2C+Nidhi)
Tichkule, S. (/jspui/browse?type=author&value=Tichkule%2C+S.)
Pandit, Shashi Bhushan (/jspui/browse?type=author&value=Pandit%2C+Shashi+Bhushan)
Chattopadhyay, K. (/jspui/browse?type=author&value=Chattopadhyay%2C+K.)

Keywords: Disulphide
C-terminal
thermostable direct
hemolysin

Issue Date: 2017

Publisher: Portland Press Ltd

Citation: Biochemical Journal, 474 (2)

Abstract: Pore-forming toxins (PFTs) are typically produced as water-soluble monomers, which upon interacting with target cells assemble into transmembrane oligomeric pores. *Vibrio parahaemolyticus* thermostable direct hemolysin (TDH) is an atypical PFT that exists as a tetramer in solution, prior to membrane binding. The TDH structure highlights a core β -sandwich domain similar to those found in the eukaryotic actinoporin family of PFTs. However, the TDH structure harbors an extended C-terminal region (CTR) that is not documented in the actinoporins. This CTR remains tethered to the β -sandwich domain through an intra-molecular disulphide bond. Part of the CTR is positioned at the interprotomer interface in the TDH tetramer. Here we show that the truncation, as well as mutation, of the CTR compromise tetrameric assembly, and the membrane-damaging activity of TDH. Our study also reveals that intra-protomer disulphide bond formation during the folding/assembly process of TDH restrains the CTR to mediate its participation in the formation of inter-protomer contact, thus facilitating TDH oligomerization. However, once tetramerization is achieved, disruption of the disulphide bond does not affect oligomeric assembly. Our study provides critical insights regarding the regulation of the oligomerization mechanism of TDH, which has not been previously documented in the PFT family. © 2017 The Author(s); published by Portland Press Limited on behalf of the Biochemical Society.

URI: <https://pubmed.ncbi.nlm.nih.gov/27784764/> (<https://pubmed.ncbi.nlm.nih.gov/27784764/>)
<http://hdl.handle.net/123456789/2705> (<http://hdl.handle.net/123456789/2705>)


Appears in Research Articles (/jspui/handle/123456789/9)
Collections:

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/2705/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/2705/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/2705?mode=full>)

 (</jspui/handle/123456789/2705/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.